Relations and Functions

Question1

Let $\mathrm{A}=\{1,2,3,4\}$ and $\mathrm{B}=\{1,4,9,16\}$. Then the number of many-one functions $f: \mathrm{A} \rightarrow \mathrm{B}$ such that $1 \in f(\mathrm{A})$ is equal to :

JEE Main 2025 (Online) 22nd January Evening Shift

Options:

A. 151

B. 139

C. 163

D. 127

Answer: A

Solution:

Step 1: Total Functions with $1 \in f(A)$

Any function $f: A \rightarrow B$ where $A=\{1,2,3,4\}$ and $B=\{1,4,9,16\}$ is defined by choosing one of the four elements of B for each element of A. Thus, the total number of functions is

$4^4=256$.

To count those functions where 1 appears at least once in the set $f(A)$, we can use the complementary counting method: subtract the functions that never use 1. If 1 is excluded, each element of A has only 3 choices (namely, $\{4,9,16\}$), so the number of such functions is

$3^4=81$.

Thus, the number of functions such that $1 \in f(A)$ is

$256-81=175$.

Step 2: Counting Many-One Functions

In this context, "many-one functions" are understood to be non-injective functions. Since an injective (one-to-one) function from A to B must be a permutation (because both sets have 4 elements), the number of one-to-one functions is

$4! = 24$.

It is important to note that every injective function $f : A \to B$ has $f(A) = B$ (a full permutation) which automatically means $1 \in f(A)$.

Thus, the number of many-one (non-injective) functions $f : A \to B$ with $1 \in f(A)$ is found by subtracting the one-to-one functions from the total functions that include 1:

$175 - 24 = 151$.

$\boxed{151}$

This detailed explanation shows that the number of many-one functions $f : A \to B$ such that $1 \in f(A)$ is indeed 151.

Question2

Let $f(x) = \log_e x$ and $g(x) = \frac{x^4-2x^3+3x^2-2x+2}{2x^2-2x+1}$. Then the domain of $f \circ g$ is

JEE Main 2025 (Online) 23rd January Morning Shift

Options:

A. $(0, \infty)$

B. $[1, \infty)$

C. $\mathbb{R}$

D. $[0, \infty)$

Answer: C

Solution:

$f(x) = \ln x$
$g(x) = \dfrac{x^4 - 2x^3 + 3x^2 - 2x + 2}{2x^2 - 2x + 1}$
$D_g \in R$
$D_f \in (0, \infty)$

For $D_{fog} \Rightarrow g(x) > 0$

$$\frac{x^4 - 2x^3 + 3x^2 - 2x + 2}{2x^2 - 2x + 1} > 0$$
$$\Rightarrow x^4 - 2x^3 + 3x^2 - 2x + 2 > 0$$

Clearly x < 0 satisfies which are included in option (1) only.

Question3

Let $f(x) = \frac{2^{x+2}+16}{2^{2x+1}+2^{x+4}+32}$. Then the value of $8\left(f\left(\frac{1}{15}\right) + f\left(\frac{2}{15}\right) + \ldots + f\left(\frac{59}{15}\right)\right)$ is equal to

JEE Main 2025 (Online) 24th January Morning Shift

Options:

A. 108

B. 92

C. 118

D. 102

Answer: C

Solution:

$$f(x) = \frac{42^x + 16}{2.2^{2x} + 16.2^x + 32}$$
$$f(x) = \frac{2\left(2^x + 4\right)}{2^{2x} + 8.2^x + 16}$$
$$f(x) = \frac{2}{2^x + 4}$$
$$f(4 - x) = \frac{2^x}{2\left(2^x + 4\right)}$$
$$f(x) + f(4 - x) = \frac{1}{2}$$

So, $\quad \mathrm{f}\left(\frac{1}{15}\right) + \mathrm{f}\left(\frac{59}{15}\right) = \frac{1}{2}$

Similarly $= f\left(\frac{29}{15}\right) + f\left(\frac{31}{15}\right) = \frac{1}{2}$

$f\left(\frac{30}{15}\right) = f(2) = \frac{2}{2^2+4} = \frac{2}{8} = \frac{1}{4}$

$\Rightarrow 8\left(29 \times \frac{1}{2} + \frac{1}{4}\right)$

Ans. 118

Question4

The function $f : (-\infty, \infty) \to (-\infty, 1)$, defined by $f(x) = \frac{2^x - 2^{-x}}{2^x + 2^{-x}}$ is :

JEE Main 2025 (Online) 24th January Evening Shift

Options:

A. One-one but not onto

B. Onto but not one-one

C. Both one-one and onto

D. Neither one-one nor onto

Answer: A

Solution:

$f(x) = \frac{2^{2x} - 1}{2^{2x} + 1}$

$= 1 - \frac{2}{2^{2x} + 1}$

$f'(x) = \frac{2}{(2^{2x}+1)^2} \cdot 2 \cdot 2^{2x} \cdot \ln 2$ i.e always $+$ ve

so $f(x)$ is $\uparrow$ function

$\therefore f(-\infty) = -1$

$f(\infty) = 1$

$\therefore f(x) \in (-1, 1) \neq$ co-domain

so function is one-one but not onto

Question5

Let $f : \mathbb{R} \to \mathbb{R}$ be a function defined by $f(x) = (2 + 3a)x^2 + \left(\frac{a+2}{a-1}\right)x + b, a \neq 1$. If $f(x + y) = f(x) + f(\mathrm{y}) + 1 - \frac{2}{7}x\mathrm{y}$, then the value of $28 \sum_{i=1}^{5} |f(i)|$ is

JEE Main 2025 (Online) 28th January Morning Shift

Options:

A. 735

B. 675

C. 715

D. 545

Answer: B

Solution:

$$f(x) = (3a + 2)x^2 + \left(\frac{a + 2}{a - 1}\right)x + b$$

$$f\left(x + \frac{1}{2}\right) = f(x) + f(y) + 1 - \frac{2}{7}xy \quad \text{.... (1)}$$

In (1) Put $x = y = 0 \Rightarrow f(0) = 2f(0) + 1 \Rightarrow f(0) = -1$

So, $\mathrm{f}(0) = 0 + 0 + \mathrm{b} = -1 \Rightarrow \mathrm{b} = -1$

In (1) Put $y = -x \Rightarrow f(0) = f(x) + f(-x) + 1 + \frac{2}{7}x^2$

$$-1 = 2(3a + 2)x^2 + 2b + 1 + \frac{2}{7}x^2$$

$$-1 = \left(2(3a + 2) + \frac{2}{7}\right)x^2 + 1 - 2$$

$$\Rightarrow 6a + 4 + \frac{2}{7} = 0$$

$$a = -\frac{5}{7}$$

So $f(x) = -\frac{1}{7}x^2 - \frac{3}{4}x - 1$

$\Rightarrow |f(x)| = \frac{1}{28}\left|4x^2 + 21x + 28\right|$

Now, $28\sum_{i=1}^{5}|f(i)| = 28(|f(1)| + |f(2)| + \ldots + |f(5)|)$

$28 \cdot \frac{1}{28} \cdot 675 = 675$

Question6

If $f(x) = \frac{2^x}{2^x+\sqrt{2}}, x \in \mathbb{R}$, then $\sum_{k=1}^{81} f\left(\frac{k}{82}\right)$ is equal to

JEE Main 2025 (Online) 28th January Morning Shift

Options:

A. 82

B. $81\sqrt{2}$

C. 41

D. $\frac{81}{2}$

Answer: D

Solution:

$$f(x) = \frac{2^x}{2^x + \sqrt{2}}$$

$$f(x) + f(1-x) = \frac{2^x}{2^x + \sqrt{2}} + \frac{2^{1-x}}{2^{1-x} + \sqrt{2}}$$

$$= \frac{2^x}{2^x + \sqrt{2}} + \frac{2}{2 + \sqrt{2}2^x} = \frac{2^x + \sqrt{2}}{2^x + \sqrt{2}} = 1$$

$$\text{Now, } \sum_{k=1}^{81} f\left(\frac{k}{82}\right) = f\left(\frac{1}{82}\right) + f\left(\frac{2}{82}\right) + \ldots\ldots + f\left(\frac{81}{82}\right)$$

$$= f\left(\frac{1}{82}\right) + f\left(\frac{2}{82}\right) + \ldots\ldots + f\left(1 - \frac{2}{82}\right) + f\left(1 - \frac{1}{82}\right)$$

$$= \left[f\left(\frac{1}{82}\right) + f\left(1 - \frac{1}{82}\right)\right] + \left[f\left(\frac{2}{82}\right) + f\left(1 - \frac{2}{82}\right)\right] + \ldots\ldots \text{40 cases } + f\left(\frac{41}{82}\right)$$

$= (1+1+\ldots 40 \text{ times }) + \frac{2^{1/2}}{2^{1/2}+2^{1/2}}$

$40 + \frac{1}{2} = \frac{81}{2}$

Question7

Let $f : [0,3] \to$ A be defined by $f(x) = 2x^3 - 15x^2 + 36x + 7$ and $g : [0,\infty) \to B$ be defined by $g(x) = \frac{x^{2025}}{x^{2025}+1}$, If both the functions are onto and $S = \{x \in Z; x \in A$ or $x \in B\}$, then $n(S)$ is equal to :

JEE Main 2025 (Online) 28th January Evening Shift

Options:

A.

29

B.

31

C.

30

D.

36

Answer: C

Solution:

as $f(x)$ is onto hence A is range of $f(x)$

now $f'(x) = 6x^2 - 30x + 36$
$= 6(x-2)(x-3)$

$f(2) = 16 - 60 + 72 + 7 = 35$

$f(3) = 54 - 135 + 108 + 7 = 34$
$f(0) = 7$

hence range $\in [7, 35] = A$

also for range of $g(x)$

$g(x) = 1 - \frac{1}{x^{2025}+1} \in [0,1) = B$
$s = \{0, 7, 8, \ldots..35\}$ hence $n(s) = 30$

Question8

If the domain of the function $\log_5(18x - x^2 - 77)$ is (α, β) and the domain of the function $\log_{(x-1)}\left(\frac{2x^2+3x-2}{x^2-3x-4}\right)$ is (γ, δ), then $\alpha^2 + \beta^2 + \gamma^2$ is equal to:

JEE Main 2025 (Online) 29th January Evening Shift

Options:

A.

186

B.

179

C.

195

D.

174

Answer: A

Solution:

$f_1(x) = \log_5(18x - x^2 - 77)$

$\therefore 18x - x^2 - 77 > 0$

$x^2 - 18x + 77 < 0$

$x \in (7, 11) \alpha = 7, \beta = 11$

$f_2(x) = \log_{(x-1)}\left(\frac{2x^2 + 3x - 2}{x^2 - 3x - 4}\right)$

$\therefore \quad x - 1 > 0, x - 1 \neq 1, \frac{2x^2 + 3x - 2}{x^2 - 3x - 4} > 0$

$x > 1, x \neq 2, \frac{(2x - 1)(x + 2)}{(x - 4)(x + 1)} > 0$

$x > 1, x \neq 2,$



$\therefore \quad x \in (4, \infty)$

$\therefore \gamma = 4$

$\therefore \alpha^2 + \beta^2 + \gamma^2 = 49 + 121 + 16$

$= 186$

Question9

If the domain of the function $f(x) = \frac{1}{\sqrt{10+3x-x^2}} + \frac{1}{\sqrt{x+|x|}}$ is (a, b), then $(1 + a)^2 + b^2$ is equal to :

JEE Main 2025 (Online) 2nd April Evening Shift

Options:

A. 29

B. 30

C. 25

D. 26

Answer: D

Solution:

$x + |x| = \begin{cases} 2x, & x \geq 0 \\ 0, & x < 0 \end{cases}$

$\Rightarrow \frac{1}{\sqrt{x+|x|}}$, domain is $x > 0$, as $2x \neq 0$

Similarly,

$\frac{1}{\sqrt{3x+10-x^2}}$ is defined when $3x + 10 - x^2 > 0$
$\Rightarrow x^2 - 3x - 10 < 0$
$(x-5)(x+2) < 0$
$\Rightarrow x \in (-2, 5)$
$\Rightarrow$ Domain will be $(0, \infty) \cap (-2, 5) = (0, 5)$
$\Rightarrow (1+a)^2 + b^2 = 1 + 25 = 26$

Question10

If the domain of the function is $f(x) = \log_e\left(\frac{2x-3}{5+4x}\right) + \sin^{-1}\left(\frac{4+3x}{2-x}\right)$ is $[\alpha, \beta)$, then $\alpha^2 + 4\beta$ equal to

Options:

A. 4

B. 3

C. 7

D. 5

Answer: A

Solution:

$\frac{2x-3}{4x+5} > 0$



$\therefore \quad x \in \left(-\infty, -\frac{5}{4}\right) \cup \left(\frac{3}{2}, \infty\right) \ldots\ldots\ldots (i)$

$-1 \leq \frac{3x+4}{2-x} \leq 1$

$\frac{3x+4}{2-x} \leq 1$

$\Rightarrow \quad \frac{3x+4}{2-x} - 1 \leq 0$

$\Rightarrow \quad \frac{3x+4-2+x}{x-2} \geq 0$

$\Rightarrow \quad \frac{4x+2}{x-2} \geq 0$

$\Rightarrow \quad x \in \left(-\infty, -\frac{1}{2}\right] \cup (2, \infty) \ldots\ldots\ldots (ii)$

$\frac{3x+4}{2-x} \geq -1$

$\Rightarrow \frac{3x+4}{2-x} + 1 \geq 0$

$\Rightarrow \frac{3x+4+2-x}{2-x} \geq 0$

$\Rightarrow \frac{2x+6}{x-2} \leq 0$

$\therefore x \in [-3, 2) \ldots\ldots\ldots (iii)$

Taking intersection of (i), (ii) and (iii)

$x \in \left[-3, -\frac{5}{4}\right)$

$\alpha = -3, \beta = -\frac{5}{4}$

$\alpha^2 + 4\beta = 4$

Question11

If the domain of the function $f(x) = \log_7\left(1 - \log_4\left(x^2 - 9x + 18\right)\right)$ is $(\alpha, \beta) \cup (\gamma, o)$, then $\alpha + \beta + \gamma + \hat{o}$ is equal to

JEE Main 2025 (Online) 3rd April Evening Shift

Options:

A. 17

B. 15

C. 16

D. 18

Answer: D

Solution:

$1-\log_4\left(x^2-9x+18\right)>0$
$\log_4\left(x^2-9x+18\right)<1$
$x^2-9x+18<4$
$x^2-9x+14<0$
$x\in(2,7)$
$x^2-9x+18>0$
$x\in(-\infty,3)\cup(6,\infty)$



$x\in(2,3)\cup(6,7)$
$\alpha+\beta+\gamma+\delta=18$

Question12

Let f be a function such that $f(x)+3f\left(\frac{24}{x}\right)=4x, x\neq 0$. Then $f(3)+f(8)$ is equal to

JEE Main 2025 (Online) 3rd April Evening Shift

Options:

A. 13

B. 11

C. 10

D. 12

Answer: B

Solution:

$f(x)+3f\left(\frac{24}{x}\right)=4x, x\neq 0 \quad \ldots(1)$

replace x by $\frac{24}{x}$

$f\left(\frac{24}{x}\right)+3f\left(\frac{24}{24}\right)=4\left(\frac{24}{x}\right)=\frac{96}{x} \quad \ldots(2)$

$3\times(2)-(1)$

$\Rightarrow 8f(x)=\frac{96.3}{x}-4x \Rightarrow f(x)=\frac{36}{x}-\frac{x}{2}$

$f(3)+f(8)=\left(12-\frac{3}{2}\right)+\left(\frac{36}{8}-4\right)$

$=8+\frac{36}{8}-\frac{12}{8}=11$

Question13

Let $f, g:(1, \infty) \rightarrow \mathbb{R}$ be defined as $f(x)=\frac{2x+3}{5x+2}$ and $g(x)=\frac{2-3x}{1-x}$. If the range of the function fog: $[2,4] \rightarrow \mathbb{R}$ is $[\alpha, \beta]$, then $\frac{1}{\beta-\alpha}$ is equal to

JEE Main 2025 (Online) 4th April Morning Shift

Options:

A. 56

B. 2

C. 29

D. 68

Answer: A

Solution:

$g(2)=4, g(4)=\frac{10}{3}$

f is decreasing in $\left(\frac{10}{3}, 4\right)$

$\therefore \quad \alpha=f(4)=\frac{1}{2}$

$\beta=f\left(\frac{10}{3}\right)=\frac{29}{56}$

$\frac{1}{\beta-\alpha}=\frac{1}{\frac{29}{56}-\frac{1}{2}}=56$

Question14

Let the domains of the functions $f(x)=\log_4 \log_3 \log_7\left(8-\log_2\left(x^2+4x+5\right)\right)$ and $g(x)=\sin^{-1}\left(\frac{7x+10}{x-2}\right)$ be (α, β) and $[\gamma, \delta]$, respectively. Then $\alpha^2+\beta^2+\gamma^2+\delta^2$ is equal to :

JEE Main 2025 (Online) 4th April Evening Shift

Options:

A. 15

B. 13

C. 16

D. 14

Answer: A

Solution:

$f(x)=\log_4\left(\log_3\left(\log_7\left(8-\log_2\left(x^2+4x+5\right)\right)\right)\right)$

$\log_3\left(\log_1\left(8-\log_2\left(x^2+4x+5\right)\right)\right)>0$

$\log_7\left(8-\log_2\left(x^2+4x+5\right)\right)>1$

$8-\log_2\left(x^2+4x+5\right)>7$

$-\log_2\left(x^2+4x+5\right)>-1$

$\log_2\left(x^2+4x+5\right)<1$

$x^2+4x+5<2$

$x^2+4x+3<0$

$\Rightarrow(x+3)(x+1)<0 \quad \ldots(1)$

$\log_7\left(8-\log_2\left(x^2+4x+5\right)\right)>0$

$8-\log_2\left(x^2+4x+5\right)>1$

$\log_2\left(x^2+4x+5\right)<9$

$x^2+4x+5<2^9$

$x^2+4x+5<512$

$\Rightarrow x^2+4x-507<0$

$\Rightarrow x=-4\pm\sqrt{16+2028}$

$x=\dfrac{-4\pm\sqrt{2044}}{2} \quad \ldots\ldots(2)$

$\Rightarrow\left(x-\left(\dfrac{-4+\sqrt{2044}}{2}\right)\right)\left(x-\left(\dfrac{-4-\sqrt{2044}}{2}\right)\right)<0$

$x^2+4x+5>0$

$D>0$

$x\in R$

Also, $8-\log_2\left(x^2+4x+5\right)>0$

$\log_2\left(x^2+4x+5\right)<8$

$x^2+4x+5<256$

$\Rightarrow x^2+4x-251<0$

$\Rightarrow x=-4\pm\sqrt{16+1004}$

$\Rightarrow x=\dfrac{-4\pm\sqrt{1020}}{2}$

$\Rightarrow\left(x-\left(\dfrac{-4+\sqrt{1020}}{2}\right)\right)\left(x-\left(\dfrac{-4-\sqrt{1020}}{2}\right)\right)<0$

$\therefore$ Intersection of (1), (2) and (3)



$\therefore x\in(-3,-1)$

$-1\le\dfrac{7x+10}{x-2}\le1$

$\Rightarrow x\in[-2,-1]$

$\therefore\alpha^2+\beta^2+\gamma^2+\delta^2=(-3)^2+(-1)^2+(-2)^{-2}+(-1)^2$

$=9+1+4+1$

$=15$

Question15

If the range of the function $f(x)=\frac{5-x}{x^2-3x+2},\ x\neq 1,2$, is $(-\infty,\alpha]\cup[\beta,\infty)$, then $\alpha^2+\beta^2$ is equal to :

JEE Main 2025 (Online) 7th April Evening Shift

Options:

A.

188

B.

192

C.

190

D.

194

Answer: D

Solution:

$$y=\frac{5-x}{x^2-3x+2}$$
$$yx^2-3xy+2y+x-5=0$$
$$yz^2+(-3y+1)x+(2y-5)=0$$

Case I : If $y=0$ (Accepted)

$\Rightarrow x=5$

Case II : If $y\neq 0$

$D \geq 0$
$(-3y+1)^2 - 4(y)(2y-5) \geq 0$
$9y^2 + 1 - 6y - 8y^2 + 20y \geq 0$
$y^2 + 14y + 1 \geq 0$
$(y+7)^2 - 48 \geq 0$
$|y+7| \geq 4\sqrt{3}$
$\Rightarrow y+7 \geq 4\sqrt{3}$ or $y+7 \leq -4\sqrt{3}$
$\Rightarrow y \geq 4\sqrt{3}-7$ or $y \leq -4\sqrt{3}-7$

From Case I and Case II

$y \in (-\infty, -4\sqrt{3}-7] \cup [4\sqrt{3}-7, \infty)$

So $\alpha = -4\sqrt{3}-7$
$\beta = 4\sqrt{3}-7$
$\Rightarrow a^2 + b^2 = (-4\sqrt{3}-7)^2 + (4\sqrt{3}-7)^2$
$= 2(48+49)$
$= 194$

Question16

Let the domain of the function $f(x) = \cos^{-1}\left(\frac{4x+5}{3x-7}\right)$ be $[\alpha, \beta]$ and the domain of $g(x) = \log_2\left(2 - 6\log_{27}(2x+5)\right)$ be (γ, δ).

Then $|7(\alpha+\beta) + 4(\gamma+\delta)|$ is equal to _____.

JEE Main 2025 (Online) 8th April Evening Shift

Answer: 96

Solution:

$f(x) = \cos^{-1}\left(\frac{4x+5}{3x-7}\right)$

$\Rightarrow -1 \leq \left(\frac{4x+5}{3x-7}\right) \leq 1$

$\left(\frac{4x+5}{3x-7}\right) \geq -1$

$\frac{4x+5+3x-7}{3x-7} \geq 0$

$\Rightarrow \frac{7x-2}{3x-7} \geq 0$



$x \in \left(-\infty, \frac{2}{7}\right] \cup \left(\frac{7}{3}, \infty\right)$

$\& \frac{4x+5}{3x-7} \leq 1 \Rightarrow \frac{x+12}{3x-7} \leq 0$



$\therefore$ Domain of f(x) is

$[-12, \frac{2}{7}] \alpha = -12, \beta = \frac{2}{7}$

$g(x) = \log_2(2 - 6\log_{27}(2x+5))$

Domain

$2 - 6\log_{27}(2x+5) > 0$

$\Rightarrow \quad 6\log_{27}(2x+5) < 2$

$\Rightarrow \quad \log_{27}(2x+5) < \frac{1}{3}$

$\Rightarrow \quad 2x+5 < 3$

$\Rightarrow \quad x < -1$

$\& 2x+5 > 0 \Rightarrow x > -\frac{5}{2}$

Domain is $x \in \left(-\frac{5}{2}, -1\right)$

$\gamma = -\frac{5}{2}, \delta = -1$

$|7(\alpha+\beta)+4(\gamma+\delta)| = \left|7\left(-12+\frac{2}{7}+4\left(-\frac{5}{2}-1\right)\right|\right.$

$|-82-14| = 96$

Question17

The function f : N− {1} → N ; defined by f(n) = the highest prime factor of n, is :

[27-Jan-2024 Shift 1]

Options:

A.

both one-one and onto

B.

one-one only

C.

onto only

D.

neither one-one nor onto

Answer: D

Solution:

$f : N - \{1\} \rightarrow N$

f(n) = The highest prime factor of n.

f(2) = 2

f(4) = 2

$\Rightarrow$ many one

4 is not image of any element

$\Rightarrow$ into

Hence many one and into

Neither one-one nor onto.

Question18

Let $f : R - \left\{ \frac{-1}{2} \right\} \rightarrow R$ and $g : R - \left\{ \frac{-5}{2} \right\} \rightarrow R$ be defined as $f(x) = \frac{2x+3}{2x+1}$ and $g(x) = \frac{|x|+1}{2x+5}$. Then the domain of the function fog is :

[27-Jan-2024 Shift 2]

Options:

A.

$$R - \left\{ -\frac{5}{2} \right\}$$

B.

R

C.

$$R - \left\{ -\frac{7}{4} \right\}$$

D.

$$R - \left\{ -\frac{5}{2}, -\frac{7}{4} \right\}$$

Answer: A

Solution:

$$f(x) = \frac{2x+3}{2x+1}; x \neq -\frac{1}{2}$$

$$g(x) = \frac{|x|+1}{2x+5}, x \neq -\frac{5}{2}$$

Domain of f(g(x))

$$f(g(x)) = \frac{2g(x)+3}{2g(x)+1}$$

$$x \neq -\frac{5}{2} \text{ and } \frac{|x|+1}{2x+5} \neq -\frac{1}{2}$$

$$x \in R - \left\{-\frac{5}{2}\right\} \text{ and } x \in R$$

$$\therefore \text{ Domain will be } R - \left\{-\frac{5}{2}\right\}$$

Question19

Consider the function f : [1/2, 1] → R defined by $f(x) = 4\sqrt{2}x 3^{-3}\sqrt{2}x^{-1}$. Consider the statements

(I) The curve y = f(x) intersects the x-axis exactly at one point

(II) The curve y = f(x) intersects the x-axis at x = cos π/12

Then

[29-Jan-2024 Shift 1]

Options:

A.

Only (II) is correct

B.

Both (I) and (II) are incorrect

C.

Only (I) is correct

D.

Both (I) and (II) are correct

Answer: D

Solution:

$f'(x) = 12\sqrt{2}x^2 - 3\sqrt{2} \geq 0$ for $\left[\frac{1}{2}, 1\right]$

$f\left(\frac{1}{2}\right) < 0$

$f(1) > 0 \Rightarrow$ (A) is correct.

$f(x) = \sqrt{2}(4x^3 - 3x) - 1 = 0$

Let $\cos\alpha = x$,

$\cos 3\alpha = \cos\frac{\pi}{4} \Rightarrow \alpha = \frac{\pi}{12}$

$x = \cos\frac{\pi}{12}$

(4) is correct.

Question20

If $f(x) = \begin{cases} 2+2x & -1 \leq x < 0 \\ 1 - \frac{x}{3} & 0 \leq x \leq 3 \end{cases}$

$g(x) = \begin{cases} -x & -3 \leq x \leq 0 \\ x & 0 < x \leq 1 \end{cases}$ ..,

then range of (fog(x)) is

[29-Jan-2024 Shift 1]

Options:

A.

(0, 1]

B.

[0, 3)

C.

[0, 1]

D.

[0, 1)

Answer: C

Solution:

$$f(g(x)) = \begin{cases} 2+2g(x), & -1 \le g(x) < 0 \ldots\ldots (1) \\ 1-\frac{g(x)}{3}, & 0 \le g(x) \le 3 \ldots\ldots\ldots (2) \end{cases}$$

By (1) $x \in \phi$

And by (2) $x \in [-3, 0]$ and $x \in [0, 1]$



Range of $f(g(x))$ is [0, 1]

Question21

Let $f(x) = 2^x - x^2$, $x \in R$. If m and n are respectively the number of points at which the curves $y = f(x)$ and $y = f'(x)$ intersects the x-axis, then the value of m + n is

[29-Jan-2024 Shift 1]

Answer: 5

Solution:



$\therefore m = 3$

$f'(x) = 2^x \ln 2 - 2x = 0$

$2^x \ln 2 = 2x$



$\therefore n = 2$

$\Rightarrow m + n = 5$

Question22

If the domain of the function $f(x) = \cos^{-1}(2 - |x|/4) + (\log_e(3 - x))^{-1}$ is $[-\alpha, \beta) - \{y\}$, then $\alpha + \beta + \gamma$ is equal to :

[30-Jan-2024 Shift 1]

Options:

A.

12

B.

9

C.

11

D.

8

Answer: C

Solution:

$$-1 \le \left| \frac{2-|x|}{4} \right| \le 1$$

$$\Rightarrow \left| \frac{2-|x|}{4} \right| \le 1$$

$$-4 \le 2-|x| \le 4$$

$$-6 \le -|x| \le 2$$

$$-2 \le |x| \le 6$$

$$|x| \le 6$$

$$\Rightarrow x \in [-6, 6] \text{ .....(1)}$$

Now, $3-x \neq 1$

And $x \neq 2$ .....(2)

and $3-x > 0$

$x < 3$ .....(3)

From (1), (2) and (3)

$$\Rightarrow x \in [-6, 3) - \{2\}$$

$\alpha = 6$

$\beta = 3$

$\gamma = 2$

$\alpha + \beta + \gamma = 11$

Question23

Let A = {1, 2, 3,....7} and let P(1) denote the power set of A. If the number of functions f : A → P(A) such that a ∈ f(a), ∀ a ∈ A is mn,m and n∈N and m is least, then m + n is equal to_____

[30-Jan-2024 Shift 1]

Answer: 44

Solution:

$f : A \longrightarrow P(A)$

$a \in f(a)$

That means 'a' will connect with subset which contain element ' a '.

Total options for 1 will be 2^6 . (Because 2^6 subsets contains 1)

Similarly, for every other element

Hence, total is $2^6 \times 2^6 \times 2^6 \times 2^6 \times 2^6 \times 2^6 \times 2^6 = 2^{42}$

Ans. $2 + 42 = 44$

Question24

If the domain of the function f(x) = $\log_e \left(\frac{2x+3}{4x^2+x-3} \right) + \cos^{-1} \left(\frac{2x-1}{x+2} \right)$ is (α, β], then the value of 5β − 4α is equal to

[30-Jan-2024 Shift 2]

Options:

A.

10

B.

12

C.

11

D.

9

Answer: B

Solution:

$$\frac{2x+3}{4x^2+x-3}>0 \text{ and } -1\le \frac{2x-1}{x+2}\le 1$$

$$\frac{2\times+3}{(4x-3)(x+1)}>0 \quad \frac{3x+1}{x+2}\ge 0 \ \& \ \frac{x-3}{x+2}\le 0$$



$(-\infty,-2)\cup\left[\frac{-1}{3},\infty\right)$ ..... (1)

$(-2,3]$ ..... (2)

$\left[\frac{-1}{3},3\right]$ ..... (3) (1) ∩ (2) ∩ (3)

$\left(\frac{3}{4},3\right]$

$\alpha=\frac{3}{4}\beta=3$

$5\beta-4\alpha=15-3=12$

Question25

Let f : R → R be a function defined f(x) = $\frac{x}{(1+x^4)^{1/4}}$ and g(x) = f(f(f(f(x)))) then 18 $\int_0^{\sqrt{2\sqrt{5}}} x^2 g(x)\,dx$

[30-Jan-2024 Shift 2]

Options:

A.

33

B.

36

C.

42

D.

39

Answer: D

Solution:

$$f(x) = \frac{x}{(1+x^4)^{1/4}}$$

$$\text{fof}(x) = \frac{f(x)}{(1+f(x)^4)^{1/4}} = \frac{\frac{x}{(1+x^4)^{1/4}}}{\left(1+\frac{x^4}{1+x^4}\right)^{1/4}} = \frac{x}{(1+2x^4)^{1/4}}$$

$$f(f(f(f(x)))) = \frac{x}{(1+4x^4)^{1/4}}$$

$$18\int_0^{\sqrt{2\sqrt{5}}} \frac{x^3}{(1+4x^4)^{1/4}}\,dx$$

Let $1+4x^4 = t^4$

$16x^3\,dx = 4t^3\,dt$

$$\frac{18}{4}\int_1^3 \frac{t^3\,dt}{t}$$

$$= \frac{9}{2}\left(\frac{t^3}{3}\right)_1^3$$

$$= \frac{3}{2}[26] = 39$$

Question26

If $f(x)=\frac{4x+3}{6x-4}, x \neq \frac{2}{3}$ and (fof) $(x)=g(x)$, where $g: \mathbb{R}-\left\{\frac{2}{3}\right\} \rightarrow \mathbb{R}-\left\{\frac{2}{3}\right\}$, then (gogog) (4) is equal to

[31-Jan-2024 Shift 1]

Options:

A.

$$-\frac{19}{20}$$

B.

19/20

C.

-4

D.

4

Answer: D

Solution:

$$f(x)=\frac{4x+3}{6x-4}$$

$$g(x)=\frac{4\left(\frac{4x+3}{6x-4}\right)+3}{6\left(\frac{4x+3}{6x-4}\right)-4}=\frac{34x}{34}=x$$

$$g(x)=x \therefore g(g(g(4)))=4$$

Question27

If the function f : (−∞, −1] ⟶ (a, b] defined by f(x) = e^{x^3-3x+1} is one-one and onto, then the distance of the point

P(2b + 4, a+ 2) from the line $x + e^{-3}y = 4$ is :

[31-Jan-2024 Shift 2]

Options:

A.

$2\sqrt{1+e^6}$

B.

$4\sqrt{1+e^6}$

C.

$3\sqrt{1+e^6}$

D.

$\sqrt{1+e^6}$

Answer: A

Solution:

$f(x) = e^{x^3 - 3x + 1}$

$f'(x) = e^{x^3 - 3x + 1} \cdot (3x^2 - 3)$

$= e^{x^3 - 3x + 1} \cdot 3(x - 1)(x + 1)$

For $f'(x) \geq 0$

$\therefore f(x)$ is increasing function

$\therefore a = e^{-\infty} = 0 = f(-\infty)$

$b = e^{-1+3+1} = e^3 = f(-1)$

$P(2b + 4, a + 2)$

$\therefore P(2e^3 + 4, 2)$



$$d = \frac{(2e^3 + 4) + 2e^{-3} - 4}{\sqrt{1 + e^{-6}}} = 2\sqrt{1 + e^6}$$

Question28

Let $f : R \longrightarrow R$ and $g : R \longrightarrow R$ be defined as

$$f(x) = \begin{cases} \log_e x & , \quad x > 0 \\ e^{-x} & , \quad x \leq 0 \end{cases}.$$

and

$$g(x) = \begin{cases} x & , \quad x \geq 0 \\ e^x & , \quad x < 0 \end{cases}.$$

Then, gof : $R \longrightarrow R$ is :

[1-Feb-2024 Shift 1]

Options:

A.

one-one but not onto

B.

neither one-one nor onto

C.

onto but not one-one

D.

both one-one and onto

Answer: B

Solution:

$$g(f(x)) = \begin{cases} f(x) & f(x) \geq 0 \\ e^{f(x)} & f(x) < 0 \end{cases}.$$

$$g(f(x)) = \begin{cases} e^{-x} & (-\infty, 0] \\ e^{\ln x} & (0, 1) \\ \ln x & [1, \infty) \end{cases}.$$



Graph of $g(f(x))$

$g(f(x)) \Rightarrow$ Many one into

Question29

If the domain of the function f(x) = $\frac{\sqrt{x^2-25}}{(4-x^2)}+\log_{10}(x^2+2x-15)$ is $(-\infty, \alpha)U[\beta, \infty)$, then $\alpha^2 + \beta^3$ is equal to :

[1-Feb-2024 Shift 2]

Options:

A.

140

B.

175

C.

150

D.

125

Answer: C

Solution:

$$f(x) = \frac{\sqrt{x^2-25}}{4-x^2}+\log_{10}(x^2+2x-15)$$

Domain $: x^2-25 \geq 0 \Rightarrow x \in (-\infty, -5] \cup [5, \infty)$

$4-x^2 \neq 0 \Rightarrow x \neq \{-2, 2\}$

$x^2+2x-15 > 0 \Rightarrow (x+5)(x-3) > 0$

$\Rightarrow x \in (-\infty, -5) \cup (3, \infty)$

$\therefore x \in (-\infty, -5) \cup [5, \infty)$

$\alpha = -5; \beta = 5$

$\therefore \alpha^2+\beta^3 = 150$

Question30

Let A = {1, 2, 3, 4, ..., 10} and B = {0, 1, 2, 3, 4}. The number of elements in the relation $R = \{(a, b) \in A \times A: 2(a-b)^2 + 3(a-b) \in B\}$ is _____.
[6-Apr-2023 shift 1]

Answer: 18

Solution:

Solution:
$A = \{1, 2, 3, \ldots\ldots 10\}$
$B = \{0, 1, 2, 3, 4\}$
$R = \{(a, b) \in A \times A : 2(a-b)^2 + 3(a-b) \in B\}$
Now $2(a-b)^2 + 3(a-b) = (a-b)(2(a-b)+3)$
$\Rightarrow a = b$ or $a - b = -2$
When $a = b \Rightarrow 10$ order pairs
When $a - b = -2 \Rightarrow 8$ order pairs
Total = 18

Question31

Let A = {0, 34, 6, 7, 8, 9, 10} and R be the relation defined on A such that $R = \{x, y) \in A \times A : x - y$ is odd positive integer or $x - y = 2\}$. The minimum number of elements that must be aadded to the relation R, so that it is a symmetric relation, is equal to _____.
[8-Apr-2023 shift 1]

Answer: 19

Solution:

Solution:
$A = \{0, 3, 4, 6, 7, 8, 9, 10\}$ $3, 7, 9 \rightarrow$ odd
$R = \{x - y =$ odd $+$ ve or $x - y = 2\}$ $0, 4, 6, 8, 10 \rightarrow$ even
${}^{3}C_{1} \cdot {}^{5}C_{1} = 15 + (6, 4), (8, 6), (10, 8), (9, 7)$
Min ^{m} ordered pairs to be added must be
$:15 + 4 = 19$

Question32

Let A = {1, 2, 3, 4, 5, 6, 7}. Then the relation $R = \{(x, y) \in A \times A : x + y = 7\}$ is [8-Apr-2023 shift 2]

Options:

A. Symmetric but neither reflexive nor transitive

B. Transitive but neither symmetric nor reflexive

C. An equivalence relation

D. Reflexive but neither symmetric nor transitive

Answer: A

Solution:

Solution:
$R = \{(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)\}$

Question33

Let A = {2, 3, 4} and B = {8, 9, 12}. Then the number of elements in the relation $R = \{((a_1, b_1), (a_2, b_2)) \in (A \times . B, A \times B) : a_1$ divides b_2 and a_2 divides $b_1\}$ is [10-Apr-2023 shift 2]

Options:

A. 18

B. 24

C. 12

D. 36

Answer: D

Solution:

Solution:



a_1 divides b_2
Each elements has 2 choices
$\Rightarrow 3 \times 2 = 6$
a_2 divides b_1
Each elements has 2 choices
$\Rightarrow 3 \times 2 = 6$
Total $= 6 \times 6 = 36$

Question34

Let A = {1, 3, 4, 6, 9} and B = {2, 4, 5, 8, 10}. Let R be a relation defined on A × B such that R = { ((a_1, b_1), (a_2, b_2..)): $a_1 \leq b_2$ and $b_1 \leq a_2$ }. Then the number of elements in the set R is [11-Apr-2023 shift 2]

Options:

A. 52

B. 160

C. 26

D. 180

Answer: B

Solution:

Solution:
Let $a_1 = 1 \Rightarrow 5$ choices of b_2
$a_1 = 3 \Rightarrow 4$ choices of b_2
$a_1 = 4 \Rightarrow 4$ choices of b_2
$a_1 = 6 \Rightarrow 2$ choices of b_2
$a_1 = 9 \Rightarrow 1$ choices of b_2
For $(a_1, b_2)16$ ways .
Similarly, $b_1 = 2 \Rightarrow 4$ choices of a_2
$b_1 = 4 \Rightarrow 3$ choices of a_2
$b_1 = 5 \Rightarrow 2$ choices of a_2
$b_1 = 8 \Rightarrow 1$ choices of a_2
Required elements in $R = 160$

Question35

The number of the relations, on the set {1, 2, 3} containing (1, 2) and (2, 3), which are reflexive and transitive but not symmetric, is _____.
[12-Apr-2023 shift 1]

Answer: 3

Solution:

Solution:
$A = \{1, 2, 3\}$
For Reflexive $(1, 1)(2, 2), (3, 3) \in R$
For transitive : $(1, 2)$ and $(2, 3) \in R \Rightarrow (1, 3) \in R$
Not symmetric : $(2, 1)$ and $(3, 2) \notin R$
$R_1 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)\}$
$R_2 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)(2, 1)\}$
$R_3 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3), (1, 3)(2, 1)\}$

Question36

Let A = {−4, −3, −2, 0, 1, 3, 4} and R = { (a, b) ∈ A × A : b = | a | . or ^{2}=a + 1 } be a relation on A. Then the minimum number of elements, that must be added to the relation R so that it becomes reflexive and symmetric, is _____
[13-Apr-2023 shift 2]

Answer: 7

Solution:

Solution:
R = [(−4, 4), (−3, 3), (3, −2), (0, 1), (0, 0), (1, 1),
(4, 4), (3, 3) }
For reflexive, add ⇒(−2, −2), (−4, −4), (−3, −3)
For symmetric, add ⇒(4, −4), (3, −3), (−2, 3), (1, 0)

Question37

Let A = {1, 2, 3, 4} and R be a relation on the set A × A defined by R = { ((a, b, (c, d) : 2a + 3b = 4c + 5d }. Then the number of elements in R is _____
[15-Apr-2023 shift 1]

Answer: 6

Solution:

Solution:
A = {1, 2, 3, 4}
R = {(a, b), (c, d)}
2a + 3b = 4c + 5d = α let
2a = {2, 4, 6, 8} 4c = {4, 8, 12, 16}
3b = {3, 6, 9, 12} 5d = {5, 10, 15, 20}

$2a+3b=\left\{\begin{matrix}5 & 8 & 11 & 14\\ 7 & 10 & 13 & 16\\ 9 & 12 & 15 & 18\\ 11 & 14 & 17 & 20\end{matrix}\right\}$ $4c+5d\left\{\begin{matrix}9 & 14 & 19 & 24\\ 13 & 18\ldots & & \\ 17 & 22\ldots. & & \\ 21 & 26\ldots. & & \end{matrix}\right\}$

Possible value of $\alpha = 9, 13, 14, 14, 17, 18$

Pairs of $\{(a, b), (c, d)\} = 6$

Question38

Let $5f(x)+4f\left(\frac{1}{x}\right)=\frac{1}{x}+3,\ x>0$. Then $18\int_1^2 f(x)\,dx$ is equal to : [6-Apr-2023 shift 1]

Options:

A. $10\log_e 2-6$

B. $10\log_e 2+6$

C. $5\log_e 2-3$

D. $5\log_e 2+3$

Answer: A

Solution:

Solution:

$5f(x)+4f\left(\frac{1}{x}\right)=\frac{1}{x}+3\ldots(1)$

$x\to\frac{1}{x}$

$5f\left(\frac{1}{x}\right)+4f(x)=x+3\ldots(2)$

$(1)\times 5-(2)\times 4$

$\Rightarrow f(x)=\frac{5}{9x}-\frac{4}{9}x+\frac{1}{3}$

$\Rightarrow 18\int_1^2 f(x)\,dx=18\left(\frac{5}{9}\ln 2-\frac{4}{9}\times\frac{3}{2}+\frac{1}{3}\right)$

$=10\ln 2-6$

Question39

Let $A = \{x \in \mathbb{R} : [x+3] + [x+4] \le 3\}$,

$B = \left\{ x \in \mathbb{R} : 3^x \left(\sum_{r=1}^{\infty} \frac{3}{10^x} \right)^{x-3} < 3^{-3x} \right\}$, where [t] denotes greatest integer function. Then,
[6-Apr-2023 shift 1]

Options:

A. $A \subset B, A \neq B$

B. $A \cap B = \varphi$

C. $A = B$

D. $B \subset C, A \neq B$

Answer: C

Solution:

Solution:
$A = \{x \in \mathbb{R} : [x+3] + [x+4] \le 3\}$
$2[x] + 7 \le 3$
$2[x] \le -4$
$[x] \le -2 \Rightarrow x < -1$ . . . (A)

$B = \left\{ x \in \mathbb{R} : 3^x \left(\sum_{r=1}^{\infty} \frac{3}{10^x} \right)^{x-3} < 3^{-3x} \right\}$

$3^x \left(\sum_{r-1}^{\infty} \frac{3}{10^x} \right)^{x-3} < 3^{-3x}$

$3^{2x-3} \left(\frac{10}{10} \right)^{x-3}$

$\Rightarrow \left(\frac{1}{9} \right)^{x-3} < 3^{-5x}$
$\Rightarrow 3^{6-2x} < 3^{3-5x}$
$\Rightarrow 6 - 2x < 3 - 5x$
$\Rightarrow 3 < -3x$
$\Rightarrow \frac{1}{10})^x < -1$ . . . (B)
$A = B$

Question40

Let the sets A and B denote the domain and range respectively of the function f (x) = $\frac{1}{\sqrt{[x]-x}}$, where [x] denotes the smallest integer greater than or equal to x. Then among the statements :
(S1) : A ∩ B = (1, ∞) − N and
(S2) : A ∪ B = (1, ∞)
[6-Apr-2023 shift 2]

Options:

A. only (S1) is true

B. neither (S1) nor (S2) is true

C. only (S2) is true

D. both (S1) and (S2) are true

Answer: A

Solution:

Solution:

$f(x) = \frac{1}{\sqrt{[x]-x}}$

If $x \in I$ $[x] = [x]$ (greatest integer function)

If $x \notin I [x] = [x] + 1$

$$\Rightarrow f(x) = \begin{cases} \frac{1}{\sqrt{[x]-x}} & x \in I \\ \frac{1}{\sqrt{[x]+1-x}} & x \notin I \end{cases}$$

$$\Rightarrow f(x) = \begin{cases} \frac{1}{\sqrt{-\{x\}}} & x \in I \quad (\text{ does not exist }) \\ \frac{1}{\sqrt{1-\{x\}}} & x \notin I \end{cases}.$$

$\Rightarrow$ domain of $f(x) = R - I$

Now, $f(x) = \frac{1}{\sqrt{1-\{x\}}}, x \notin I$

$\Rightarrow x < \{x\} < 1$

$\Rightarrow 0 < 1\sqrt{1-\{x\}} < 1$

$\Rightarrow \frac{1}{\sqrt{1-\{x\}}} > 1$

$\Rightarrow$ Range$(1, \infty)$

$\Rightarrow A = R - I$

$B = (1\infty)$
So, $A \cap B = (1, \infty) - N$
$A \cup B \neq (1, \infty)$
$\Rightarrow$ S1 is only correct.

Question41

Let $f, g : \mathbb{N} - \{1\} \rightarrow \mathbb{N}$ be functions defined by $f(a) = \alpha$, where α is the maximum of the powers of those primes p such that p^{α} divides a, and $g(a) = a + 1$, for all $a \in \mathbb{N} - \{1\}$. Then, the function $f + g$ is [27-Jul-2022-Shift-1]

Options:

A. one-one but not onto

B. onto but not one-one

C. both one-one and onto

D. neither one-one nor onto

Answer: D

Solution:

Solution:
$f, g : N - \{1\} \rightarrow N$ defined as
$f(a) = \alpha$, where α is the maximum power of those primes p such that p^{α} divides a.
$g(a) = a + 1$
Now,
$f(2) = 1, \quad g(2) = 3 \quad \Rightarrow \quad (f + g)(2) = 4$
$f(3) = 1, \quad g(3) = 4 \quad \Rightarrow \quad (f + g)(3) = 5$
$f(4) = 2, \quad g(4) = 5 \Rightarrow (f + g)(4) = 7$
$f(5) = 1, \quad g(5) = 6 \Rightarrow (f + g)(5) = 7$
$\because (f + g)(5) = (f + g)(4)$
$\therefore f + g$ is not one-one
Now, $\because f_{min} = 1, g_{min} = 3$
So, there does not exist any $x \in N - \{1\}$ such that $(f + g)(x) = 1, 2, 3$
$\therefore f + g$ is not onto

Question42

If domain of the function $\log_e\left(\frac{6x^2+5x+1}{2x-1}\right)+\cos^{-1}\left(\frac{2x^2-3x+4}{3x-5}\right)$ is $(\alpha, \beta) \cup (\gamma, \delta]$, then, $18(\alpha^2+\beta^2+\gamma^2+\delta^2)$ is equal to
[8-Apr-2023 shift 2]

Answer: 20

Solution:

Solution:

$$\frac{6x^2+5x+1}{2x-1} > 0$$

$$\frac{(3x+1)(2x+1)}{2x-1} > 0$$

$$x \in \left[\frac{-1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right] \cup \left(\frac{5}{3}, \infty\right) \ldots \text{(B)}$$

$$x < \frac{5}{3} \ldots \text{(C)}$$

$$A \cap B \cap C \equiv \left(\frac{-1}{2}, \frac{-1}{3}\right) \cup \left(\frac{1}{2}, \frac{1}{\sqrt{2}}\right]$$

$$\text{So } 18(\alpha^2+\beta^2+\gamma^2+\delta^2) = 18\left(\frac{1}{4}+\frac{1}{9}+\frac{1}{4}+\frac{1}{2}\right)$$

$$= 18+2 = 20$$

Question43

Let R = {a, b, c, d, e} and S = {1, 2, 3, 4}. Total number of onto functions f : R → S such that f(a) ≠ 1, is equal to _____.
[8-Apr-2023 shift 2]

Answer: 180

Solution:

Solution:
Total onto function
$\frac{\sqcup 5}{\lfloor 3 \lfloor} \times \lfloor 4 = 240$
Now when $f(a) = 1$
$\left\lfloor 4 + \frac{\lfloor 4}{\lfloor 2 \lfloor 2} \times \lfloor x3 = 24 + 36 = 60.$
so required $f^{n} = 240 - 60 = 180$

Question44

If the domain of the function f(x) = $\sec^{-1}\left(\frac{2x}{5x+3}\right)$ is [α, β)U(γ, δ], then $|3\alpha + 10(\beta + \gamma) + 21\delta|$ is equal to _____.
[10-Apr-2023 shift 2]

Answer: 24

Solution:

Solution:
$f(x) = \sec^{-1}\frac{2x}{5x+3}$
$\left|\frac{2x}{5x+3}\right|$
$\left|\frac{2x}{5x+3}\right| \geq 1 \Rightarrow |2x| \geq |5x+3|$
$(2x)^2 - (5x+3)^2 \geq 0$
$(7x+3)(-3x-3) \geq 0$



$\therefore$ domain $\left[-1, \frac{-3}{5}\right) \cup \left(\frac{-3}{5}, \frac{-3}{7}\right]$
$\alpha = -1, \beta = \frac{-3}{5}, \gamma = \frac{-3}{5}, \delta = \frac{-3}{7}$
$3\alpha + 10(\beta + \gamma) + 21\delta = -3$
$-3 + 10\left(\frac{-6}{5}\right) + \left(\frac{-3}{7}\right)21 = -24$

Question45

The domain of the function f (x) = $\frac{1}{\sqrt{[x]^2-3[x]-10}}$ is (where [x] denotes the greatest integer less than or equal to x)
[11-Apr-2023 shift 2]

Options:

A. $(-\infty, -3] \cup [6, \infty)$

B. $(-\infty, -2) \cup (5, \infty)$

C. $(-\infty, -3] \cup (5, \infty)$

D. $(-\infty, -2) \cup [6, \infty)$

Answer: D

Solution:

Solution:

$$F(x) = \frac{1}{\sqrt{[x]^2-3[x]-10}}$$

$[x]^2 - 3[x] - 10 > 0$

$([x] + 2)([x] - 5) > 0$

$[x] < -2$ or $[x] > 5$

$[x] \leq -3$ or $[x] \geq 6$

$x < -2$ or $x \geq 6$

$x \in (-\infty, -2) \cup [6, \infty)$

Question46

Let A = {1, 2, 3, 4, 5} and B = {1, 2, 3, 4, 5, 6}. Then the number of functions f : A → B satisfying f (1) + f (2) = f(4) − 1 is equal to _____.
[11-Apr-2023 shift 2]

Answer: 360

Solution:

Solution:
$f(1)+f(2)+1=f(4)\leq 6$
$f(1)+f(2)\leq 5$
Case (i) $f(1)=1 \Rightarrow f(2)=1,2,3,4 \Rightarrow 4$ mappings
Case (ii) $f(1)=2 \Rightarrow f(2)=1,2,3 \Rightarrow 3$ mappings
Case (iii) $f(1)=3 \Rightarrow f(2)=1,2 \Rightarrow 2$ mappings
Case (iv) $f(1)4 \Rightarrow f(2)=1 \Rightarrow 1$ mapping
$f(5)\&f(6)$ both have 6 mappings each
Number of functions $=(4+3+2+1)\times 6\times 6=360$

Question47

Let D be the domain of the function $f(x)=\sin^{-1}\left(\log_{3x}\left(\frac{6+2\log_3 x}{-5x}\right)\right)$. If the range of the function $g: D \rightarrow R$ defined by $g(x)=x-[x]$, ($[x]$ is the greatest integer function), is (α, β), then $\alpha^2+\frac{5}{\beta}$ is equal to [12-Apr-2023 shift 1]

Options:

A. 46

B. 135

C. 136

D. 45

Answer: B

Solution:

Solution:

$\frac{6+2\log_3 x}{-5x} > 0 \& x > 0 \& x \neq \frac{1}{3}$

this gives $x \in \left(0, \frac{1}{27}\right) \ldots (1)$

$-1 \leq \log_{3x}\left(\frac{6+2\log_3 x}{-5x}\right) \leq 1$

$3x \leq \frac{6+2\log_3 x}{-5x} \leq \frac{1}{3x}$

$15x^2 + 6 + 2\log_3 x \geq 0 \quad 6 + 2\log_3 x + \frac{5}{3} \geq 0$

$x \in \left(0, \frac{1}{27}\right) \ldots (2) \quad x \geq 3^{-\frac{23}{6}} \ldots . (3)$

form (1), (2) & (3)

$x \in \left[3^{-\frac{23}{6}}, \frac{1}{27}\right)$

$\therefore \alpha$ is small positive quantity

$\& \beta = \frac{1}{27}$

$\therefore \alpha^2 + \frac{5}{\beta}$ is just greater than 135

Question48

For x ∈ R, two real valued functions f(x) and g(x) are such that, $g(x) = \sqrt{x} + 1$ and $fog(x) = x + 3 - \sqrt{x}$. Then f(0) is equal to [13-Apr-2023 shift 1]

Options:

A. 5

B. 0

C. -3

D. 1

Answer: A

Solution:

Solution:

$g(x) = \sqrt{x} + 1$

$fog\ (x) = x + 3 - \sqrt{x}$

$= (\sqrt{x} + 1)^2 - 3(\sqrt{x} + 1) + 5$

$= g^2(x) - 3g(x) + 5$

$\Rightarrow f(x) = x^2 - 3x + 5$

$\therefore f(0) = 5$

But, if we consider the domain of the composite function fog (x) then in that case $f(0)$ will be not defined as $g(x)$ cannot be equal to zero.

Question49

For the differentiable function $f : R - \{0\} \rightarrow R$, let $3f(x) + 2f\left(\frac{1}{x}\right) = \frac{1}{x} - 10$, then $\left|f(3) + f'\left(\frac{1}{4}\right)\right|$ is equal to [13-Apr-2023 shift 1]

Options:

A. 13

B. $\frac{29}{5}$

C. $\frac{33}{5}$

D. 7

Answer: A

Solution:

Solution:

$\left[3f(x) + 2f\left(\frac{1}{x}\right) = \frac{1}{x} - 10\right] \times 3$

$\left[2f(x) + 3f\left(\frac{1}{x}\right) = x - 10\right] \times 2$

$5f(x) = \frac{3}{x} - 2x - 10$

$f(x) = \frac{1}{5}\left(\frac{3}{x} - 2x - 10\right)$

$f'(x) = \frac{1}{5}\left(-\frac{3}{x^2} - 2\right)$

$\left|f(3) + f'\left(\frac{1}{4}\right)\right| = \left|\frac{1}{5}(1 - 6 - 10) + \frac{1}{5}(-48 - 2)\right|$

$= |-3 - 10| = 13$

Question50

The range of f (x) = $4\sin^{-1}\left(\frac{x^2}{x^2+1}\right)$ is

[13-Apr-2023 shift 2]

Options:

A. [0, π)

B. [0, π]

C. [0, 2π)

D. [0, 2π]

Answer: C

Solution:

Solution:

$f(x) = 4\sin^{-1}\left(\frac{x^2}{1+x^2}\right)$

$0 \le \frac{x^2}{1+x^2} < 1$

$\Rightarrow 0 \le \sin^{-1}\left(\frac{x^2}{1+x^2}\right) < \frac{\pi}{2}$

$\Rightarrow 0 \le 4\sin^{-1}\left(\frac{x^2}{1+x^2}\right) < 2\pi$

Range : [0, 2π)

Question51

If the domain of the function

$f(x) = \log_e(4x^2+11x+6) + \sin^{-1}(4x+3) + \cos^{-1}\left(\frac{10x+6}{3}\right)$ is (α, β], then

36 | α + β| is equal to

[15-Apr-2023 shift 1]

Options:

A. 72

B. 63

C. 45

D. 54

Answer: C

Solution:

Solution:

$f(x) = \ln(4x^2 + 11x + 6) + \sin^{-1}(4x + 3)$

$+ \cos^{-1}\left(\frac{10x + 6}{3}\right)$

(i) $4x^2 + 11x + 6 > 0$

$4x^2 + 8x + 3x + 6 > 0$

$(4x + 3)(x + 2) > 0$

$x \in (-\infty, -2) \cup \left(-\frac{3}{4}, \infty\right)$

(ii) $4x + 3 \in [-1, 1]$

$x \in [-1, -1/2]$

(iii) $\frac{10x + 6}{3} \in [-1, 1]$

$x \in \left[-\frac{9}{10}, -\frac{3}{10}\right]$

$x \in \left(-\frac{3}{4}, -\frac{1}{2}\right] \quad \alpha = -\frac{3}{4}, \beta = -\frac{1}{2}$

$\alpha + \beta = -\frac{5}{4}$

$36 | \alpha + \beta | = 45$

Question52

The relation R = {(a, b) : gcd(a, b) = 1, 2a ≠ b, a, b ∈ ℤ} is: [24-Jan-2023 Shift 1]

Options:

A. transitive but not reflexive

B. symmetric but not transitive

C. reflexive but not symmetric

D. neither symmetric nor transitive

Answer: D

Solution:

Solution:
Reflexive : (a, a) gcd of $(a, a) = 1$
Which is not true for every a $\in Z$.
Symmetric:
Take $a = 2, b = 1$ $\gcd(2, 1) = 1$
Also $2a = 4 \neq b$
Now when $a = 1, b = 2$ $\gcd(1, 2) = 1$
Also now $2a = 2 = b$
Hence $a = 2b$
$\Rightarrow R$ is not Symmetric
Transitive:
Let $a = 14, b = 19, c = 21$
$\gcd(a, b) = 1$
$\gcd(b, c) = 1$
$\gcd(a, c) = 7$
Hence not transitive
R is neither symmetric nor transitive.

Question53

Let R be a relation defined on N as a R b is 2a + 3b is a multiple of 5, a, b ∈ ℕ. Then R is
[29-Jan-2023 Shift 2]

Options:

A. not reflexive

B. transitive but not symmetric

C. symmetric but not transitive

D. an equivalence relation

Answer: D

Solution:

Solution:
a $Ra \Rightarrow 5a$ is multiple it 5
So reflexive

$aRb \Rightarrow 2a + 3b = 5\alpha,$
Now b R a

$$2b + 3a = 2b + \left(\frac{5\alpha - 3b}{2}\right) \cdot 3$$

$$= \frac{15}{2}\alpha - \frac{5}{2}b = \frac{5}{2}(3\alpha - b)$$

$$= \frac{5}{2}(2a + 2b - 2\alpha)$$

$$= 5(a + b - \alpha)$$

Hence symmetric

$a\ R\ b \Rightarrow 2a + 3b = 5\alpha$.

$b\ R\ c \Rightarrow 2b + 3c = 5\beta$

Now $2a + 5b + 3c = 5(\alpha + \beta)$

$\Rightarrow 2a + 5b + 3c = 5(\alpha + \beta)$

$\Rightarrow 2a + 3c = 5(\alpha + \beta - b)$

$\Rightarrow aRc$

Hence relation is equivalence relation.

Question54

The minimum number of elements that must be added to the relation R = {(a, b), (b, c)} on the set {a, b, c} so that it becomes symmetric and transitive is:
[30-Jan-2023 Shift 1]

Options:

A. 4

B. 7

C. 5

D. 3

Answer: B

Solution:

Solution:
For Symmetric $(a, b), (b, c) \in R$
$\Rightarrow (b, a), (c, b) \in R$
For Transitive $(a, b), (b, c) \in R$
$\Rightarrow (a, c) \in R$
Now
1. Symmetric
$\therefore (a, c) \in R \Rightarrow (c, a) \in R$

2. Transitive
$\therefore (a, b), (b, a) \in R$
$\Rightarrow (a, a) \in R \& (b, c), (c, b) \in R$
$\Rightarrow (b, b) \& (c, c) \in R$
$\therefore$ Elements to be added

$$\{(b, a) \quad (c, b) \quad (a, c) \quad (c, a), (a, a) \quad (b, b) \quad (c, c)\}$$

Number of elements to be added $= 7$

Question55

Let R be a relation on N × N defined by (a, b)R (c, d) if and only if ad (b − c) = bc(a − d). Then R is
[31-Jan-2023 Shift 1]

Options:

A. symmetric but neither reflexive nor transitive

B. transitive but neither reflexive nor symmetric

C. reflexive and symmetric but not transitive

D. symmetric and transitive but not reflexive

Answer: A

Solution:

Solution:
$(a, b)R(c, d) \Rightarrow ad(b - c) = bc(a - d)$
Symmetric:
$(c, d)R(a, b) \Rightarrow cb(d - a) = da(c - b) \Rightarrow$
Symmetric
Reflexive:
$(a, b)\ R(a, b) \Rightarrow ab(b - a) \neq ba(a - b) \Rightarrow$
Not reflexive
Transitive: $(2, 3)R(3, 2)$ and $(3, 2)R(5, 30)$ but
$((2, 3), (5, 30)) \notin R \Rightarrow$ Not transitive

Question56

Among the relations

$S = \left\{ (a, b) : a, b \in \mathbb{R} - \{0\}, 2 + \frac{a}{b} > 0 \right\}$

And $T = \{(a, b) : a, b \in \mathbb{R}, a^2 - b^2 \in Z\}$,
[31-Jan-2023 Shift 2]

Options:

A. S is transitive but T is not

B. T is symmetric but S is not

C. Neither S nor T is transitive

D. Both S and T are symmetric

Answer: B

Solution:

Solution:
For relation $T = a^2 - b^2 = -I$
Then, (b, a) on relation R
$\Rightarrow b^2 - a^2 = -I$
$\therefore T$ is symmetric

$S = \left\{ (a, b) : a, b \in R - \{0\}, 2 + \frac{a}{b} > 0 \right\}$

$2 + \frac{a}{b} > 0 \Rightarrow \frac{a}{b} > -2, \Rightarrow \frac{b}{a} < \frac{-1}{2}$

If $(b, a) \in S$ then

$2 + \frac{b}{a}$ not necessarily positive

$\therefore S$ is not symmetric

Question57

Let R be a relation on $\mathbb{R}$, given by $R = \{(a, b) : 3a - 3b + \sqrt{7}$ is an irrational number $\}$. Then R is
[1-Feb-2023 Shift 1]

Options:

A. Reflexive but neither symmetric nor transitive

B. Reflexive and transitive but not symmetric

C. Reflexive and symmetric but not transitive

D. An equivalence relation

Answer: A

Solution:

Solution:
Check for reflexivity:
As 3(a-a) $+\sqrt{7} = \sqrt{7}$ which belongs to relation so relation is reflexive
Check for symmetric:
Take $a = \frac{\sqrt{7}}{3}, b = 0$
Now $(a, b) \in R$ but $(b, a) \notin R$
As $3(b - a) + \sqrt{7} = 0$ which is rational so relation is not symmetric.
Check for Transitivity:
Take (a, b) as $\left(\frac{\sqrt{7}}{3}, 1\right)$
&(b, c) as $\left(1, \frac{2\sqrt{7}}{3}\right)$
So now $(a, b) \in R$&$(b, c) \in R$ but $(a, c) \notin R$ which means relation is not transitive

---

Question58

Let P(S) denote the power set of S = {1, 2, 3, ..., 10}. Define the relations R_1 and R_2 on P(S) as AR_1B if $(A \cap B^c) \cup (B \cap A^9) = \emptyset$ and AR_2B if $A \cup B^c = B \cup A^c$, $\forall A, B \in P(S)$. Then : [1-Feb-2023 Shift 2]

Options:

A. both R_1 and R_2 are equivalence relations

B. only R_1 is an equivalence relation

C. only R_2 is an equivalence relation

D. both R_1 and R_2 are not equivalence relations

Answer: A

Solution:

Solution:
S = {1, 2, 3, … . .10}
P(S) = power set of S
AR, B $\Rightarrow (A \cap \overrightarrow{B}) \cup (\overrightarrow{A} \cap B) = \varphi$
R1 is reflexive, symmetric
For transitive
$(A \cap \overrightarrow{B}) \cup (\overrightarrow{A} \cap B) = \varphi$; {a} = φ = {b} A = B
$(B \cap \overrightarrow{C}) \cup (\overrightarrow{B} \cap C) = \varphi \therefore B = C$
$\therefore A = C$ equivalence.



$R_2 \equiv A \cup \overrightarrow{B} = \overrightarrow{A} \cup B$
$R_2 \rightarrow$ Reflexive, symmetric
for transitive



$A \cup \overrightarrow{B} = \overrightarrow{A} \cup B \Rightarrow$ {a, c, d} = {b, c, d}
{a} = {b} $\therefore A = B$
$B \cup \overrightarrow{C} = \overrightarrow{B} \cup C \Rightarrow B = C$
$\therefore A = C \quad \therefore A \cup \overrightarrow{C} = \overrightarrow{A} \cup C \therefore$ Equivalence

---

Question59

The equation $x^2 - 4x + [x] + 3 = x[x]$, where [x] denotes the greatest integer function, has:
[24-Jan-2023 Shift 1]

Options:

A. exactly two solutions in $(-\infty, \infty)$

B. no solution

C. a unique solution in $(-\infty, 1)$

D. a unique solution in $(-\infty, \infty)$

Answer: D

Solution:

Solution:

$x^2 - 4x + [x] + 3 = x[x]$

$\Rightarrow x^2 - 4x + 3 = x[x] - [x]$

$(x-1)(x-3) = [x] \cdot (x-1)$

$\Rightarrow x = 1$ or $x - 3 = [x]$

$\Rightarrow x - [x] = 3$

$\{x\} = 3$ (Not Possible)

Only one solution $x = 1$ in $(-\infty, \infty)$

Question60

Let f (x) be a function such that f (x + y) = f (x) · f (y) for all x, y ∈ N. If f(1) = 3 and $\sum_{k=1}^{n} f(k) = 3279$, then the value of n is [24-Jan-2023 Shift 2]

Options:

A. 6

B. 8

C. 7

D. 9

Answer: C

Solution:

Solution:

$f(x + y) = f(x) \cdot f(y)\ \forall x, y \in N, f(1) = 3$

$f(2) = f^2(1) = 3^2$

$f(3) = f(1)f(2) = 3^3$

$f(4) = 3^4$

$f(k) = 3^k$

$\sum_{k=1}^{n} f(k) = 3279$

$f(1) + f(2) + f(3) + \ldots\ldots\ldots + f(k) = 3279$

$3 + 3^2 + 3^3 + \ldots\ldots\ldots . 3^k = 3279$

$\frac{3(3^k-1)}{3-1}=3279$

$\frac{3^k-1}{2}=1093$

$3^k-1=2186$

$3^k=2187$

$k=7$

Question61

If $f(x)=\frac{2^{2x}}{2^{2x}+2}$, $x \in R$ then $f\left(\frac{1}{2023}\right)+f\left(\frac{2}{2023}\right)+.....+f\left(\frac{2022}{2023}\right)$ equal to
[24-Jan-2023 Shift 2]

Options:

A. 2011

B. 1010

C. 2010

D. 1011

Answer: D

Solution:

Solution:

$f(x)=\frac{4^x}{4^x+2}$

$f(x)+f(1-x)=\frac{4^x}{4^x+2}+\frac{4^{1-x}}{4^{1-x}+2}$

$=\frac{4^x}{4^x+2}+\frac{4}{4+2(4^x)}$

$=\frac{4^x}{4^x+2}+\frac{2}{2+4^x}$

$=1$

$\Rightarrow f(x)+f(1-x)=1$

Now $f\left(\frac{1}{2023}\right)+f\left(\frac{2}{2023}\right)+f\left(\frac{3}{2023}\right)+.....+$

$.....+f\left(1-\frac{3}{2023}\right)+f\left(1-\frac{2}{2023}\right)+f\left(1-\frac{1}{2023}\right)$

Now sum of terms equidistant from beginning and end is 1

Sum $= 1 + 1 + 1 + \ldots\ldots\ldots + 1$ (1011 times)
$= 1011$

Question62

For some a, b, c $\in \mathbb{N}$, let f (x) = ax − 3 and g(x) = x^b + c, x $\in \mathbb{R}$. If (fog)$^{-1}$(x) = $\left(\frac{x-7}{2}\right)^{1/3}$ then (fog) (ac) + (gof) (b) is equal to _____.
[25-Jan-2023 Shift 1]

Answer: 2039

Solution:

Solution:
Let fog (x) = h(x)

$\Rightarrow h^{-1}(x) = \left(\frac{x-7}{2}\right)^{\frac{1}{3}}$

$\Rightarrow h(x) = fog(x) = 2x^3 + 7$

$fog\ (x) = a(x^b + c) - 3$

$\Rightarrow a = 2, b = 3, c = 5$

$\Rightarrow$ fog (ac) = fog(10) = 2007

$g(f(x) = (2x - 3)^3 + 5.$

$\Rightarrow gof(b) = gof(3) = 32$

$\Rightarrow$ sum = 2039

Question63

Let f : $\mathbb{R} \to \mathbb{R}$ be a function defined by f (x) = $\log_{\sqrt{m}}\{\sqrt{2}(\sin x - \cos x) + m - 2\}$, for some m, such that the range of f is [0, 2]. Then the value of m is _____
[25-Jan-2023 Shift 2]

Options:

A. 5

B. 3

C. 2

D. 4

Answer: A

Solution:

Solution:
Since,
$-\sqrt{2} \le \sin x - \cos x \le \sqrt{2}$
$\therefore \quad -2 \le \sqrt{2}(\sin x - \cos x) \le 2$
(Assume $\sqrt{2}(\sin x - \cos x) = k$)
$-2 \le k \le 2 \quad \ldots$ (i)
$f(x) = \log_{\sqrt{m}}(k + k - 2)$
Given,
$0 \le f(x) \le 2$
$0 \le \log_{\sqrt{\min}}(k + m - 2) \le 2$
$1 \le k + m - 2 \le m$
$-m + 3 \le k \le 2 \ldots ..ii)$
From eq. (i) & (ii), we get $-m + 3 = -2$
$\Rightarrow m = 5$

Question64

The number of functions f : {1, 2, 3, 4} → {a ∈ ℤ: | a | ≤8} satisfying f(n)+ $\frac{1}{n}$f(n + 1) = 1, ∀n ∈ {1, 2, 3} is
[25-Jan-2023 Shift 2]

Options:

A. 3

B. 4

C. 1

D. 2

Answer: D

Solution:

Solution:
$f : \{1, 2, 3, 4\} \rightarrow \{a \in \mathbb{Z} : |a| \leq 8\}$
$f(n) + \frac{1}{n} f(n+1) = 1, \ \forall n \in \{1, 2, 3\}$
$f(n+1)$ must be divisible by n
$f(4) \Rightarrow -6, -3, 0, 3, 6$
$f(3) \Rightarrow -8, -6, -4, -2, 0, 2, 4, 6, 8$
$f(2) \Rightarrow -8, \ldots\ldots\ldots\ldots, 8$
$f(1) \Rightarrow -8, \ldots\ldots\ldots\ldots, 8$
$\frac{f(4)}{3}$ must be odd since $f(3)$ should be even therefore 2 solution possible.

Question65

Let $f(x) = 2x^n + \lambda$, $\lambda \in \mathbb{R}$, $n \in \mathbb{N}$, and $f(4) = 133$, $f(5) = 255$. Then the sum of all the positive integer divisors of $(f(3) - f(2))$ is [25-Jan-2023 Shift 2]

Options:

A. 61

B. 60

C. 58

D. 59

Answer: B

Solution:

Solution:
$f(x) = 2x^n + \lambda$
$f(4) = 133$
$f(5) = 255$
$133 = 2 \times 4^n + \lambda \ldots (1)$
$255 = 2 \times 5^n + \lambda \ldots (2)$
$(2) - (1)$
$122 = 2(5^n - 4^n)$
$\Rightarrow 5^n - 4^n = 61$
$\therefore n = 3 \& \lambda = 5$
Now, $f(3) - f(2) = 2(3^3 - 2^3) = 38$
Number of Divisors is $1, 2, 19, 38$; & their sum is 60

Question66

Let f : R → R be a function such that f (x) = $\frac{x^2+2x+1}{x^2+1}$. Then [29-Jan-2023 Shift 1]

Options:

A. f (x) is many-one in (−∞, −1)

B. f (x) is many-one in (1, ∞)

C. f (x) is one-one in [1, ∞) but not in (−∞, ∞)

D. f (x) is one-one in (−∞, ∞)

Answer: C

Solution:

Solution:



$$f(x) = \frac{(x+1)^2}{x^2+1} = 1 + \frac{2x}{x^2+1}$$

$$f(x) = 1 + \frac{2}{x + \frac{1}{x}}$$

Question67

The domain of f (x) = $\frac{\log_{(x+1)}(x-2)}{e^{2\log_e x}-(2x+3)}$, x ∈ R is

[29-Jan-2023 Shift 1]

Options:

A. $\mathbb{R}-\{1-3\}$

B. $(2,\infty)-\{3\}$

C. $(-1,\infty)-\{3\}$

D. $\mathbb{R}-\{3\}$

Answer: B

Solution:

Solution:
$x-2>0 \Rightarrow x>2$
$x+1>0 \Rightarrow x>-1$
$x+1\neq 1 \Rightarrow x\neq 0$ and $x>0$
Denominator
$x^2-2x-3\neq 0$
$(x-3)(x+1)\neq 0$
$x\neq -1, 3$
So Ans $(2,\infty)-\{3\}$

Question68

Consider a function f : $\mathbb{N}\rightarrow\mathbb{R}$, satisfying
f (1) + 2f (2) + 3f (3) + … + xf (x) = x(x + 1)f (x); x ≥ 2 with f (1) = 1.
Then $\frac{1}{f(2022)}+\frac{1}{f(2028)}$ is equal to
[29-Jan-2023 Shift 2]

Options:

A. 8200

B. 8000

C. 8400

D. 8100

Answer: D

Solution:

Solution:
Given for $x \geq 2$

$f(1) + 2f(2) + ... + xf(x) = x(x+1)f(x)$

replace x by $x+1$

$\Rightarrow \quad x(x+1)f(x) + (x+1)f(x+1)$

$= (x+1)(x+2)f(x+1)$

$\Rightarrow \quad \frac{x}{f(x+1)} + \frac{1}{f(x)} = \frac{(x+2)}{f(x)}$

$\Rightarrow \quad xf(x) = (x+1)f(x+1) = \frac{1}{2}, x \geq 2$

$f(2) = \frac{1}{4}, f(3) = \frac{1}{6}$

Now $f(2022) = \frac{1}{4044}$

$f(2028) = \frac{1}{4056}$

So, $\frac{1}{f(2022)} + \frac{1}{f(2028)} = 4044 + 4056 = 8100$

Question69

Suppose $f : R \rightarrow (0, \infty)$ be a differentiable function such that $5f(x + y) = f(x) \cdot f(y), \forall x, y \in R$. If $f(3) = 320$, then $\sum_{n=0}^{5} f(n)$ is equal to: [30-Jan-2023 Shift 1]

Options:

A. 6875

B. 6575

C. 6825

D. 6528

Answer: C

Solution:

Solution:
Option (3)

$5f(x + y) = f(x) \cdot f(y)$

$5f(0) = f(0)^2 \Rightarrow f(0) = 5$

$5f(x+1) = f(x) \cdot f(1)$

$\Rightarrow \quad \frac{f(x+1)}{f(x)} = \frac{f(1)}{5}$

$\Rightarrow \quad \frac{f(1)}{f(0)} \cdot \frac{f(2)}{f(1)} \cdot \frac{f(3)}{f(2)} = \left(\frac{f(1)}{5}\right)^3$

$\Rightarrow \quad \frac{320}{5} = \frac{(f(1))^3}{5^3} \Rightarrow f(1) = 20$

$\therefore \quad 5f(x+1) = 20 \cdot f(x) \Rightarrow f(x+1) = 4f(x)$

$\sum_{n=0}^{5} f(n) = 5 + 5.4 + 5.4^2 + 5.4^3 + 5.4^4 + 5.4^5$

$= \frac{5[4^6 - 1]}{3} = 6825$

Question70

Let S = {1, 2, 3, 4, 5, 6}. Then the number of one functions f : S → P(S), where P(S) denote the power set of S, such that f (n) ⊂ f (m) where n < m is _____. [30-Jan-2023 Shift 1]

Answer: 3240

Solution:

Solution:

Let $S = \{1, 2, 3, 4, 5, 6\}$, then the number of one-one functions, $f : S \cdot P(S)$, where $P(S)$ denotes the power set of S, such that $f(n) < f(m)$ where $n < m$ is

$n(S) = 6$

$P(S) = \left\{ \begin{matrix} \varphi & \{1\} & \ldots \{6\} & \{1, 2\} & \ldots \\ \{5, 6\} & \ldots & \{1, 2, 3, 4, 5, 6\} & & \end{matrix} \right\}$

-64 elements

case -1

$f(6) = S$ i.e. 1 option,

$f(5) =$ any 5 element subset A of S i.e. 6 options,

$f(4) =$ any 4 element subset B of A i.e. 5 options,

$f(3) =$ any 3 element subset C of B i.e. 4 options,

$f(2) =$ any 2 element subset D of C i.e. 3 options,

$f(1) =$ any 1 element subset E of D or empty subset i.e. 3 options,

Total functions $= 1080$

Case - 2

$f(6)=$ any 5 element subset A of S i.e. 6 options,
$f(5)=$ any 4 element subset B of A i.e. 5 options,
f' (4) = any 3 element subset C of B i.e. 4 options,
$f(3)=$ any 2 element subset D of C i.e. 3 options,
f' $(2)=$ any 1 element subset E of D i.e. 2 options,
$f(1)=$ empty subset i.e. 1 option
Total functions $=720$
Case -3
$f(6)=S$
$f(5)=$ any 4 element subset A of' S i.e. 15 options,
$f(4)=$ any 3 element subset B of A i.e. 4 options,
$f(3)=$ any 2 element subset C of B i.e. 3 options,
$f(2)=$ any 1 element subset D of C i.e. 2 options,
$f(1)=$ empty subset i.e. 1 option

Total functions $=360$
Case -4
$f(6)=S$
$f(5)=$ any 5 element subset A of S i.e. 6 options,
$f(4)=$ any 3 element subset B of A i.e. 10 options,
$f(3)=$ any 2 element subset C of B i.e. 3 options,
$f(2)=$ any 1 element subset D of C i.e. 2 options,
$f(1)=$ empty subset i.e. 1 option
Total functions $=360$
Case -5
$f(6)=S$
$f(5)=$ any 5 element subset A of S i.e. 6 options,
$f(4)=$ any 4 element subset B of A i.e. 5 options,
$f(3)=$ any 2 element subset C of B i.e. 6 options,
$f(2)=$ any 1 element subset D of C i.e. 2 options,
$f(1)=$ empty subset i.e. 1 option
Total functions $=360$
Case - 6
$f(6)=S$
$f(5)=$ any 5 element subset A of S i.e. 6 options,
$f(4)=$ any 4 element subset B of A i.e. 5 options,
$f(3)=$ any 3 element subset C of B i.e. 4 options,
$f(2)=$ any 1 element subset D of C i.e. 3 options,
$f(1)=$ empty subset i.e. 1 option
Total functions $=360$
$\therefore$ Number of such functions $=3240$

Question71

Let $f^1(x)=\frac{3x+2}{2x+3}$, $x \in R-\left\{\frac{-3}{2}\right\}$

For $n \geq 2$, define $f^n(x)=f^1 0 f^{n-1}(x)$.

If $f^5(x) = \frac{ax+b}{bx+a}$, gcd(a, b) = 1, then a + b is equal to _____.
[30-Jan-2023 Shift 1]

Answer: 3125

Solution:

Solution:

$$f^1(x) = \frac{3x+2}{2x+3}$$

$$\Rightarrow f^2(x) = \frac{13x+12}{12x+13}$$

$$\Rightarrow f^3(x) = \frac{63x+62}{62x+63}$$

$$\therefore f^5(x) = \frac{1563x+1562}{1562x+1563}$$

$$a + b = 3125$$

Question72

The range of the function $f(x) = \sqrt{3-x} + \sqrt{2+x}$ is
[30-Jan-2023 Shift 2]

Options:

A. $[\sqrt{5}, \sqrt{10}]$

B. $[2\sqrt{2}, \sqrt{11}]$

C. $[\sqrt{5}, \sqrt{13}]$

D. $[\sqrt{2}, \sqrt{7}]$

Answer: A

Solution:

Solution:

$$y^2 = 3 - x + 2 + x + 2\sqrt{(3-x)(2+x)}$$

$$= 5 + 2\sqrt{6 + x - x^2}$$

$y^2 = 5 + 2\sqrt{\frac{25}{4} - \left(x - \frac{1}{2}\right)^2}$

$y_{max} = \sqrt{5+5} = \sqrt{10}$

$y_{min} = \sqrt{5}$

Question73

Let A = {1, 2, 3, 5, 8, 9}. Then the number of possible functions f : A → A such that f(m · n) = f(m) · f(n) for every m, n ∈ A with m · n ∈ A is equal to _____.
[30-Jan-2023 Shift 2]

Answer: 432

Solution:

Solution:
$f(1) = 1$; $f(9) = f(3) \times f(3)$
i.e., $f(3) = 1$ or 3
Total function $= 1 \times 6 \times 2 \times 6 \times 6 \times 1 = 432$

Question74

If the domain of the function $f(x) = \frac{[x]}{1+x^2}$, where [x] is greatest integer ≤x, is [2, 6), then its range is
[31-Jan-2023 Shift 1]

Options:

A. $\left(\frac{5}{26}, \frac{2}{5}\right] - \left\{\frac{9}{29}, \frac{27}{109}, \frac{18}{89}, \frac{9}{53}\right\}$

B. $\left(\frac{5}{26}, \frac{2}{5}\right]$

C. $\left(\frac{5}{37}, \frac{2}{5}\right] - \left\{\frac{9}{29}, \frac{27}{109}, \frac{18}{89}, \frac{9}{53}\right\}$

D. $\left(\frac{5}{37}, \frac{2}{5}\right]$

Answer: D

Solution:

Solution:

$f(x) = \frac{2}{1+x^2}$ $x \in [2, 3)$

$f(x) = \frac{3}{1+x^2}$ $x \in [3, 4)$

$f(x) = \frac{4}{1+x^2}$ $x \in [4, 5)$

$f(x) = \frac{5}{1+x^2}$ $x \in [5, 6)$



$\left(\frac{5}{37}, \frac{2}{5}\right]$

Question75

The absolute minimum value, of the function $f(x) = x^2 - x + 1 \mid + [x^2 - x + 1]$, where [t] denotes the greatest integer function, in the interval $[-1, 2]$, is :
[31-Jan-2023 Shift 2]

Options:

A. $\frac{3}{4}$

B. $\frac{3}{2}$

C. $\frac{1}{4}$

D. $\frac{5}{4}$

Answer: A

Solution:

Solution:
$f(x) = |x^2 - x + 1| + [x^2 - x + 1]; x \in [-1, 2]$
Let $g(x) = x^2 - x + 1$
$= \left(x - \frac{1}{2}\right)^2 + \frac{3}{4}$
$\because |x^2 - x + 1|$ and $[x^2 - x + 2]$
Both have minimum value at $x = 1/2$
$\Rightarrow$ Minimum $f(x) = \frac{3}{4} + 0$
$= \frac{3}{4}$

Question76

Let $f : \mathbb{R} - \{2, 6\} \rightarrow \mathbb{R}$ be real valued function defined as $f(x) = \frac{x^2 + 2x + 1}{x^2 - 8x + 12}$. Then range of f is

[31-Jan-2023 Shift 2]

Options:

A. $\left(-\infty, -\frac{21}{4}\right] \cup [0, \infty)$

B. $\left(-\infty, -\frac{21}{4}\right) \cup (0, \infty)$

C. $\left(-\infty, -\frac{21}{4}\right] \cup \left[\frac{21}{4}, \infty\right)$

D. $\left(-\infty, -\frac{21}{4}\right] \cup [1, \infty)$

Answer: A

Solution:

Solution:

Let $y = \dfrac{x^2+2x+1}{x^2-8x+12}$

By cross multiplying

$yx^2 - 8xy + 12y - x^2 - 2x - 1 = 0$

$x^2(y-1) - x(8y+2) + (12y-1) = 0$

Case 1, $y \neq 1$

$D \geq 0$

$\Rightarrow (8y+2)^2 - 4(y-1)(12y-1) \geq 0$

$\Rightarrow y(4y+21) \geq 0$



$y \in \left(-\infty, \dfrac{-21}{4}\right] \cup [0, \infty) - \{1\}$

Case 2, $y = 1$

$x^2 + 2x + 1 = x^2 - 8x + 12$

$10x = 11$

$x = \dfrac{11}{10}$ So, y can be 1

Hence $y \in \left(-\infty, \dfrac{-21}{4}\right] \cup [0, \infty)$

Question77

Let $f : R - \{0, 1\} \to R$ be a function such that $f(x) + f\left(\dfrac{1}{1-x}\right) = 1 + x$.

Then f (2) is equal to :
[1-Feb-2023 Shift 2]

Options:

A. $\dfrac{9}{2}$

B. $\dfrac{9}{4}$

C. $\dfrac{7}{4}$

D. $\frac{7}{3}$

Answer: B

Solution:

Solution:

$f(x)+f\left(\frac{1}{1-x}\right)=1+x$

$x=2 \Rightarrow f(2)+f(-1)=3$

$x=-1 \Rightarrow f(-1)+f\left(\frac{1}{2}\right)=0 \ldots (2)$

$x=\frac{1}{2} \Rightarrow f\left(\frac{1}{2}\right)+f(2)=\frac{3}{2} \ldots (3)$

$(1)+(3)-(2) \Rightarrow 2f(2)=\frac{9}{2}$

$\therefore f(2)=\frac{9}{4}$

Question78

Let $f: \mathbb{R} \to \mathbb{R}$ be defined as $f(x) = x - 1$ and $g: \mathbb{R} - \{1, -1\} \to \mathbb{R}$ be defined as $g(x) = \frac{x^2}{x^2-1}$. Then the function fog is : [26-Jun-2022-Shift-2]

Options:

A. one-one but not onto

B. onto but not one-one

C. both one-one and onto

D. neither one-one nor onto

Answer: D

Solution:

Solution:

$f : R \to R$ defined as

$f(x) = x - 1$ and $g : R \to \{1, -1\} \to R$, $g(x) = \dfrac{x^2}{x^2 - 1}$

Now $fog(x) = \dfrac{x^2}{x^2 - 1} - 1 = \dfrac{1}{x^2 - 1}$

$\therefore$ Domain of $fog(x) = R - \{-1, 1\}$

And range of $fog(x) = (-\infty, -1] \cup (0, \infty)$

Now, $\dfrac{d}{dx}(\text{fog}(x)) = \dfrac{-1}{x^2 - 1} \cdot 2x = \dfrac{2x}{1 - x^2}$

$\therefore \dfrac{d}{dx}(f \circ g(x)) > 0$ for $\dfrac{2x}{(1 - x)(1 + x)} > 0$

$\Rightarrow \dfrac{x}{(x - 1)(x + 1)} < 0$

$\therefore x \in (-\infty, -1) \cup (0, 1)$

and $\dfrac{d}{dx}(f \circ g(x)) < 0$ for $x \in (-1, 0) \cup (1, \infty)$

$\therefore fog(x)$ is neither one-one nor onto.

Question79

Let f : R → R be a function defined by f (x) = $\dfrac{2e^{2x}}{e^{2x} + e}$.
Then

$f\left(\frac{1}{100}\right) + f\left(\frac{2}{100}\right) + f\left(\frac{3}{100}\right) + \ldots + f\left(\frac{99}{100}\right)$
is equal to____
[27-Jun-2022-Shift-1]

Answer: 99

Solution:

Solution:
Given,

$$f(x) = \frac{2e^{2x}}{e^{2x}+e}$$

$$\therefore f(1-x) = \frac{2e^{2(1-x)}}{e^{2(1-s)}+e}$$

$$= \frac{2 \cdot \frac{e^2}{e^{2x}}}{\frac{e^2}{e^{2\pi}}+e}$$

$$= \frac{2e^2}{e^2+e^{2x}\cdot e}$$

$$= \frac{2e^2}{e(e+e^{2x})}$$

$$= \frac{2e}{e+e^{2\Delta}}$$

$$\therefore f(x)+f(1-x) = \frac{2e^{2x}}{e^{2x}+e} + \frac{2e}{e^{2x}+e}$$

$$= \frac{2(e^{2x}+e)}{e^{2x}+e}$$

$$= 2 \ldots\ldots (1)$$

Now,

$$f\left(\frac{1}{100}\right)+f\left(\frac{99}{100}\right)$$

$$= f\left(\frac{1}{100}\right)+f\left(1-\frac{1}{100}\right)$$

$= 2$ [as $f(x)+f(1-x)=2$]

$$f\left(\frac{2}{100}\right)+f\left(1-\frac{2}{100}\right) = 2$$

$$\vdots$$

$$f\left(\frac{49}{100}\right)+f\left(1-\frac{49}{100}\right) = 2$$

$\therefore$ Total sum $= 49 \times 2$

Remaining term $= f\left(\frac{50}{100}\right) = f\left(\frac{1}{2}\right)$

Put $x = \frac{1}{2}$ in equation (1), we get

$$f\left(\frac{1}{2}\right)+f\left(1-\frac{1}{2}\right) = 2$$

$$\Rightarrow 2f\left(\frac{1}{2}\right) = 2$$

$\Rightarrow f\left(\frac{1}{2}\right) = 1$

$\therefore$ Sum $= 49 \times 2 + 1 = 99$

Question80

Let S = {1, 2, 3, 4, 5, 6, 7, 8, 9, 10} . Define f : S → S as

$$\mathbf{f(n) =} \begin{cases} 2n & \text{, if } n = 1,\ 2,\ 3,\ 4,\ 5 \\ 2n - 11 & \text{, if } n = 6,\ 7,\ 8,\ 9,\ 10, \end{cases}$$

Let g : S → S be a function such that $fog(n) = \begin{cases} n+1 & \text{, if n is odd} \\ n-1 & \text{, if n is even} \end{cases}$

Then g(10)g(1) + g(2) + g(3) + g(4) + g(5)) is equal to [27-Jun-2022-Shift-2]

Answer: 190

Solution:

Solution:

$$\because f(n) = \begin{cases} 2n & n = 1\ \ 2\ \ 3\ \ 4\ \ 5 \\ 2n - 11 & n = 6\ \ 7\ \ 8\ \ 9\ \ 10 \end{cases}.$$

$\therefore f(1) = 2, f(2) = 4, \ldots\ldots, f(5) = 10$

and $f(6) = 1, f(7) = 3, f(8) = 5, \ldots\ldots, f(10) = 9$

Now, $f(g(n)) = \begin{cases} n+1 & \text{if } n \text{ is odd} \\ n-1 & \text{if } n \text{ is even} \end{cases}$.

$f(g(10)) = 9 \Rightarrow g(10) = 10$

$f(g(1)) = 2 \Rightarrow g(1) = 1$

. $f(g(2)) = 1 \Rightarrow g(2) = 6$

$\therefore f(g(3)) = 4 \Rightarrow g(3) = 2$

$f(g(4)) = 3 \Rightarrow g(4) = 7$

$f(g(5)) = 6 \Rightarrow g(5) = 3$

$\therefore g(10)g(1) + g(2) + g(3) + g(4) + g(5)) = 190$

Question81

Let a function f : N → N be defined by

$$f(n) = \begin{cases} 2n & n = 2,\ 4,\ 6,\ 8,\ \ldots\ldots \\ n-1 & n = 3,\ 7,\ 11,\ 15,\ \ldots\ldots \\ \frac{n+1}{2} & n = 1,\ 5,\ 9,\ 13,\ \ldots\ldots. \end{cases}$$

then, f is
[28-Jun-2022-Shift-1]

Options:

A. one-one but not onto

B. onto but not one-one

C. neither one-one nor onto

D. one-one and onto

Answer: D

Solution:

Solution:
When $n = 1, 5, 9, 13$ then $\frac{n+1}{2}$ will give all odd numbers.
When $n = 3, 7, 11, 15\ldots\ldots$
$n - 1$ will be even but not divisible by 4
When $n = 2, 4, 6, 8\ldots$
Then $2n$ will give all multiples of 4
So range will be N.
And no two values of n give same y, so function is one-one and onto.

Question82

The probability that a randomly chosen one-one function from the set {a, b, c, d } to the set {1, 2, 3, 4, 5} satisfies $f(a) + 2f(b) - f(c) = f(d)$ is :
[28-Jun-2022-Shift-2]

Options:

A. $\frac{1}{24}$

B. $\frac{1}{40}$

C. $\frac{1}{30}$

D. $\frac{1}{20}$

Answer: D

Solution:

Solution:
Number of one-one function from $\{a, b, c, d\}$ to set $\{1, 2, 3, 4, 5\}$ is ${}^5P_4 = 120 n(s)$.
The required possible set of value (f(a), $f(b), f(c), f(d)$) such that $f(a) + 2f(b) - f(c) = f(d)$ are (5, 3, 2, 1), (5, 1, 2, 3), (4, 1, 3, 5), (3, 1, 4, 5), (5, 4, 3, 2) and (3, 4, 5, 2)
$\therefore n(E) = 6$
$\therefore$ Required probability $= \frac{n(E)}{n(S)} = \frac{6}{120} = \frac{1}{20}$

Question83

Let S = {1, 2, 3, 4}. Then the number of elements in the set $\{f : S \times S \to S : f$ is onto and $f(a, b) = f(b, a) \geq a \ \forall (a, b) \in S \times S\}$ is____
[28-Jun-2022-Shift-2]

Answer: 37

Solution:

Solution:
There are 16 ordered pairs in $S \times S$. We write all these ordered pairs in 4 sets as follows.
$A = \{(1, 1)\}$
$B = \{(1, 4), (2, 4), (3, 4)(4, 4), (4, 3), (4, 2), (4, 1)\}$
$C = \{(1, 3), (2, 3), (3, 3), (3, 2), (3, 1)\}$
$D = \{(1, 2), (2, 2), (2, 1)\}$
All elements of set B have image 4 and only element of A has image 1 .
All elements of set C have image 3 or 4 and all elements of set D have image 2 or 3 or 4 .
We will solve this question in two cases.
Case I: When no element of set C has image 3 .
Number of onto functions $= 2$ (when elements of set D have images 2 or 3)
Case II: When atleast one element of set C has image 3 .
Number of onto functions $= (2^3 - 1)(1 + 2 + 2) = 35$
Total number of functions $= 37$

Question84

The domain of the function $\cos^{-1}\left(\frac{2\sin^{-1}\left(\frac{1}{4x^2-1}\right)}{\pi}\right)$ is :

[29-Jun-2022-Shift-1]

Options:

A. $R - \left\{-\frac{1}{2}, \frac{1}{2}\right\}$

B. $(-\infty, -1] \cup [1, \infty) \cup \{0\}$

C. $\left(-\infty, \frac{-1}{2}\right) \cup \left(\frac{1}{2}, \infty\right) \cup \{0\}$

D. $\left(-\infty, \frac{-1}{\sqrt{2}}\right] \cup \left[\frac{1}{\sqrt{2}}, \infty\right) \cup \{0\}$

Answer: D

Solution:

Solution:

$$-1 \le \frac{2\sin^{-1}\left(\frac{1}{4x^2-1}\right)}{\pi} \le 1$$

$$\Rightarrow -\frac{\pi}{2} \le \sin^{-1}\left(\frac{1}{4x^2-1}\right) \le \frac{\pi}{2}$$

$$\Rightarrow -1 \le \frac{1}{4x^2-1} \le 1$$

$$\therefore \frac{1}{4x^2-1} + 1 \ge 0$$

$$\Rightarrow \frac{1+4x^2-1}{4x^2-1} \ge 0$$

$$\Rightarrow \frac{4x^2}{4x^2-1} \ge 0$$

$$\Rightarrow \frac{4x^2}{(2x+1)(2x-1)} \ge 0 \ldots\ldots (1)$$

$$\therefore x \in \left(-\alpha, -\frac{1}{2}\right) \cup \{0\} \cup \left(\frac{1}{2}, \alpha\right)$$

And $\frac{1}{4x^2-1} - 1 \le 0$

$$\Rightarrow \frac{1-4x^2+1}{4x^2-1} \le 0$$

$$\Rightarrow \frac{2-4x^2}{4x^2-1} \le 0$$

$$\Rightarrow \frac{2x^2-1}{4x^2-1} \ge 0$$

$$\Rightarrow \frac{(\sqrt{2}x+1)(\sqrt{2}x-1)}{(2x+1)(2x-1)} \ge 0$$

$$x \in \left(-\alpha, -\frac{1}{\sqrt{2}}\right) \cup \left(-\frac{1}{2}, \frac{1}{2}\right) \cup \left(\frac{1}{\sqrt{2}}, \alpha\right)$$

From (3) and (4), we get

$$\therefore x \in \left[-\alpha, -\frac{1}{\sqrt{2}}\right) \cup \left[\frac{1}{\sqrt{2}}, \alpha\right) \cup \{0\}$$

Question85

Let c, k ∈ R. If $f(x) = (c+1)x^2 + (1-c^2)x + 2k$ and $f(x+y) = f(x) + f(y) - xy$, for all x, y ∈ R, then the value of $|2(f(1) + f(2) + f(3) + \ldots\ldots + f(20))|$ is equal to____ [29-Jun-2022-Shift-1]

Answer: 3395

Solution:

Solution:
$f(x)$ is polynomial
Put $y = 1/x$ in given functional equation we get

$$f\left(x+\frac{1}{x}\right) = f(x) + f\left(\frac{1}{x}\right) - 1$$

$$\Rightarrow (c+1)\left(x+\frac{1}{x}\right)^2 + (1-c^2)\left(x+\frac{1}{x}\right) + 2K$$

$$= (c+1)x^2 + (1-c^2)x + 2K + (c+1)\frac{1}{x^2} + (1-c^2)\frac{1}{x} + 2K - 1$$

$$\Rightarrow 2(c+1) = 2K - 1 \ldots . (1)$$

and put $x = y = 0$ we get
$f(0) = 2 + f(0) - 0 \Rightarrow f(0) = 0 \Rightarrow k = 0$
$\therefore k = 0$ and $2c = -3 \Rightarrow c = -3/2$

$$f(x) = -\frac{x^2}{2} - \frac{5x}{4} = \frac{1}{4}(5x + 2x^2)$$

$$\left|2\sum_{i=1}^{20} f(i)\right| = \left|\frac{-2}{4}\left(\frac{5.20.21}{2} + \frac{2.20.21.41}{6}\right)\right|$$

$$= \left|\frac{-1}{2}(2730 + 5740)\right|$$

$$= \left|-\frac{6790}{2}\right| = 3395.$$

Question86

Let f (x) and g(x) be two real polynomials of degree 2 and 1 respectively. If $f(g(x)) = 8x^2 - 2x$ and $g(f(x)) = 4x^2 + 6x + 1$, then the value of $f(2) + g(2)$ is_____
[29-Jun-2022-Shift-2]

Answer: 18

Solution:

Solution:
$f(g(x)) = 8x^2 - 2x.$
$g(f(x)) = 4x^2 + 6x + 1.$
So, $g(x) = 2x - 1$
&$f(x) = 2x^2 + 3x + 1$
$f(2) = 8 + 6 + 1 = 15$
Ans. 18

Question87

The domain of the function

f (x) = $\frac{\cos^{-1}\left(\frac{x^2-5x+6}{x^2-9}\right)}{\log_e(x^2-3x+2)}$ **is :**

[24-Jun-2022-Shift-1]

Options:

A. $(-\infty, 1) \cup (2, \infty)$

B. $(2, \infty)$

C. $\left[-\frac{1}{2}, 1\right) \cup (2, \infty)$

D. $\left[-\frac{1}{2}, 1\right) \cup (2, \infty) - \left\{3, \frac{3+\sqrt{5}}{2}, \frac{3-\sqrt{5}}{2}\right\}$

Answer: D

Solution:

Solution:

$$-1 \le \frac{x^2-5x+6}{x^2-9} \le 1 \text{ and } x^2-3x+2>0, \neq 1$$

$$\frac{(x-3)(2x+1)}{x^2-9} \ge 0 \mid \frac{5(x-3)}{x^2-9} \ge 0$$

The solution to this inequality is

$$x \in \left[\frac{-1}{2}, \infty\right) - \{3\}$$

for $x^2-3x+2>0$ and $\neq 1$

$$x \in (-\infty, 1) \cup (2, \infty) - \left\{\frac{3-\sqrt{5}}{2}, \frac{3+\sqrt{5}}{2}\right\}$$

Combining the two solution sets (taking intersection)

$$x \in \left[-\frac{1}{2}, 1\right) \cup (2, \infty) - \left\{\frac{3-\sqrt{5}}{2}, \frac{3+\sqrt{5}}{2}\right\}$$

Question88

The number of one-one functions f : {a, b, c, d } → {0, 1, 2, ……, 10} such that 2f (a) − f (b) + 3f (c) + f (d) = 0 is [24-Jun-2022-Shift-1]

Answer: 31

Solution:

Solution:

$$-1 \le \frac{x^2-5x+6}{x^2-9} \le 1 \text{ and } x^2-3x+2>0, \ne 1$$

$$\frac{(x-3)(2x+1)}{x^2-9} \ge 0 \mid \frac{5(x-3)}{x^2-9} \ge 0$$

The solution to this inequality is

$$x \in \left[\frac{-1}{2}, \infty\right) - \{3\}$$

for $x^2-3x+2>0$ and $\ne 1$

$$x \in (-\infty, 1) \cup (2, \infty) - \left\{\frac{3-\sqrt{5}}{2}, \frac{3+\sqrt{5}}{2}\right\}$$

Combining the two solution sets (taking intersection)

$$x \in \left[-\frac{1}{2}, 1\right) \cup (2, \infty) - \left\{\frac{3-\sqrt{5}}{2}, \frac{3+\sqrt{5}}{2}\right\}$$

f(d) can't be 9 and 10 as if $f(d) = 9$ or 10 then $f(b) = 2+9 = 11$ or $f(b) = 2+10 = 12$, which is not possible as here any function's maximum value can be 10 .

$\therefore$ Total possible functions when $f(c)=0$ and $f(a)=1$ are $=7$

(2) When $f(c)=0$ and $f(a)=2$ then

$3\times 0+2\times 2+f(d)=f(b)$

$\Rightarrow 4+f(d)=f(b)$

$\therefore$ possible value of $f(d)=1, 3, 4, 5, 6$

$\therefore$ Total possible functions in this case $=5$

(3) When $f(c)=0$ and $f(a)=3$ then

$3\times 0+2\times 3+f(d)=f(b)$

$\Rightarrow 6+f(d)=f(b)$

$\therefore$ Possible value of $f(d)=1, 2, 4$

$\therefore$ Total possible functions in this case $=3$

(4) When $f(c)=0$ and $f(a)=4$ then

$3\times 0+2\times 4+f(d)=f(b)$

$\Rightarrow 8+f(d)=f(b)$

$\therefore$ Possible value of $f(d)=1, 2$

$\therefore$ Total possible functions in this case $=2$

(5) When $f(c)=0$ and $f(a)=5$ then

$3\times 0+2\times 5+f(d)=f(b)$

$\Rightarrow 10+f(d)=f(b)$

Possible value of $f(d)$ can be 0 but $f(c)$ is already zero. So, no value to $f(d)$ can satisfy.

$\therefore$ No function is possible in this case.

$\therefore$ Total possible functions when f(c)=0 and f(a)=1,2,3 and 4 are =7+5+3+2=17

Case II:

(1) When f(c)=1 and f(a)=0 then

3×1+2×0+f(d)=f(b)

⇒3+f(d)=f(b)

$\therefore$ Possible value of f(d)=2,3,4,5,6,7

$\therefore$ Total possible functions in this case =6

(2) When f(c)=1 and f(a)=2 then

$3\times1+2\times2+f(d)=f(b)$

$\Rightarrow 7+f(d)=f(b)$

$\therefore$ Possible value of f(d)=0,3

$\therefore$ Total possible functions in this case =2

(3) When f(c)=1 and f(a)=3 then

$3\times1+2\times3+f(d)=f(b)$

$\Rightarrow 9+f(d)=f(b)$

$\therefore$ Possible value of f(d)=0

$\therefore$ Total possible functions in this case =1

$\therefore$ Total possible functions when f(c)=1 and f(a)=0,2 and 3 are =6+2+1=9

Case III:

(1) When f(c)=2 and f(a)=0 then

$3\times2+2\times0+f(d)=f(b)$

$\Rightarrow 6+f(d)=f(b)$

$\therefore$ Possible values of f(d)=1,3,4

$\therefore$ Total possible functions in this case =3

(2) When f(c)=2 and f(a)=1 then,

$3\times2+2\times1+f(d)=f(b)$

$\Rightarrow 8+f(d)=f(b)$

$\therefore$ Possible values of f(d)=0

$\therefore$ Total possible function in this case =1

$\therefore$ Total possible functions when f(c)=2 and f(a)=0,1 are =3+1=4

Case IV:

(1) When f(c)=3 and f(a)=0 then

$3\times3+2\times0+f(d)=f(b)$

$\Rightarrow 9+f(d)=f(b)$

$\therefore$ Possible values of f(d)=1

$\therefore$ Total one-one functions from four cases

=17+9+4+1=31

Question89

Let R_1 and R_2 be relations on the set {1, 2, ……., 50} such that

$R_1 = \{(p, p^n) : p.$ is a prime and $n \geq 0$ is an integer $\}$ and

$R_2 = \{(p, p^n) : p.$ is a prime and $n = 0$ or $1\}$

Then, the number of elements in $R_1 - R_2$ is___

[28-Jun-2022-Shift-1]

Answer: 8

Solution:

Solution:

$R_1 - R_2 = \{(2, 2^2), (2, 2^3), (2, 2^4), (2, 2^5), (3, 3^2), (3, 3^3), (5, 5^2), (7, 7^2)\}$

So number of elements $= 8$

Question90

Let $R_1 = \{(a, b) \in N \times N : | a - b | \leq 13\}$ and

$R_2 = \{(a, b) \in N \times N : | a - b | \neq 13\}$. Then on N :

[28-Jun-2022-Shift-2]

Options:

A. Both R_1 and R_2 are equivalence relations

B. Neither R_1 nor R_2 is an equivalence relation

C. R_1 is an equivalence relation but R_2 is not

D. R_2 is an equivalence relation but R_1 is not

Answer: B

Solution:

Solution:
$R_1 = \{(a, b) \in N \times N : |a - b| \leq 13\}$ and
$R_2 = \{(a, b) \in N \times N : |a - b| \neq 13\}$
In R_1: $\because |2 - 11| = 9 \leq 13$
$\therefore$ $(2, 11) \in R_1$ and $(11, 19) \in R_1$ but $(2, 19) \notin R_1$
$\therefore$ R_1 is not transitive
Hence R_1 is not equivalence
In R_2 : $(13, 3) \in R_2$ and $(3, 26) \in R_2$ but $(13, 26) \notin R_2 (\because |13 - 26| = 13)$
$\therefore R_2$ is not transitive
Hence R_2 is not equivalence.

Question91

The probability that a relation R from {x, y} to {x, y} is both symmetric and transitive, is equal to
[29-Jun-2022-Shift-2]

Options:

A. $\frac{5}{16}$

B. $\frac{9}{16}$

C. $\frac{11}{16}$

D. $\frac{13}{16}$

Answer: A

Solution:

Solution:
Total no. of relations $= 2^{2 \times 2} = 16$
Fav. relation $= \varphi, \{(x, x)\}, \{(y, y)\}, \{(x, x)(y, y)\}$
$\{(x, x), (y, y), (x, y)(y, x)\}$
Prob. $= \frac{5}{16}$

Question92

The number of bijective functions $f : \{1, 3, 5, 7, ..., 99\} \to \{2, 4, 6, 8, \ldots 100\}$, such that $f(3) \geq f(9) \geq f(15) \geq f(21) \geq \ldots . f(99)$, is [25-Jul-2022-Shift-2]

Options:

A. ${}^{50}P_{17}$

B. ${}^{50}P_{33}$

C. $33! \times 17!$

D. $\frac{50!}{2}$

Answer: B

Solution:

Solution:
As function is one-one and onto, out of 50 elements of domain set 17 elements are following restriction $f(3) > f(9) > f(15)..... > f(99)$
So number of ways $= {}^{50}C_{17} \cdot 1.33\ !$
$= {}^{50}P_{33}$

Question93

Let $f(x)$ be a quadratic polynomial with leading coefficient 1 such that $f(0) = p$, $p \neq 0$, and $f(1) = \frac{1}{3}$. If the equations $f(x) = 0$ and $f \circ f \circ f \circ f(x) = 0$ have a common real root, then $f(-3)$ is equal to___ [25-Jul-2022-Shift-2]

Answer: 25

Solution:

Solution:
Let $f(x) = (x-\alpha)(x-\beta)$
It is given that $f(0) = p \Rightarrow \alpha\beta = p$
and $f(1) = \frac{1}{3} \Rightarrow (1-\alpha)(1-\beta) = \frac{1}{3}$
Now, let us assume that, α is the common root of $f(x) = 0$ and fofofof $(x) = 0$
fofofof $(x) = 0$
$\Rightarrow$ fofof $(0) = 0$
$\Rightarrow f\,of\,(p) = 0$
So, $f(p)$ is either α or β.
$(p-\alpha)(p-\beta) = \alpha$
$(\alpha\beta-\alpha)(\alpha\beta-\beta) = \alpha \Rightarrow (\beta-1)(\alpha-1)\beta = 1 (\because \alpha \neq 0)$
So, $\beta = 3$
$(1-\alpha)(1-3) = \frac{1}{3}$
$\alpha = \frac{7}{6}$
$f(x) = \left(x - \frac{7}{6}\right)(x-3)$
$f(-3) = \left(-3 - \frac{7}{6}\right)(3-3) = 25$

Question94

Let f : R → R be a continuous function such that f (3x) − f (x) = x. If f (8) = 7, then f (14) is equal to:
[26-Jul-2022-Shift-1]

Options:

A. 4

B. 10

C. 11

D. 16

Answer: B

Solution:

Solution:

$f(3x) - f(x) = x$..... (1)

$x \to \frac{x}{3}$

$f(x) - f\left(\frac{x}{3}\right) = \frac{x}{3}$..... (2)

Again $x \to \frac{x}{3}$

$f\left(\frac{x}{3}\right) - f\left(\frac{x}{9}\right) = \frac{x}{3^2}$.....

Similarly

$f\left(\frac{x}{3^{n-2}}\right) - f\left(\frac{x}{3^{n-1}}\right) = \frac{x}{3^{n-1}}$.... (n)

Adding all these and applying $n \to \infty$

$\lim\limits_{n \to \infty}\left(f(3x) - f\left(\frac{x}{3^{n-1}}\right)\right) = x\left(1 + \frac{1}{3} + \frac{1}{3^2} + \ldots\right)$

$f(3x) - f(0) = \frac{3x}{2}$

Putting $x = \frac{8}{3}$

$f(8) - f(0) = 4$

$\Rightarrow f(0) = 3$

Putting $x = \frac{14}{3}$

$f(14) - 3 = 7 \Rightarrow f(14) = 0$

Question95

The domain of the function $f(x) = \sin^{-1}[2x^2 - 3] + \log_2\left(\log_{\frac{1}{2}}(x^2 - 5x + 5)\right)$, where [t] is the greatest integer function, is : [27-Jul-2022-Shift-2]

Options:

A. $\left(-\sqrt{\frac{5}{2}}, \frac{5-\sqrt{5}}{2}\right)$

B. $\left(\frac{5-\sqrt{5}}{2}, \frac{5+\sqrt{5}}{2}\right)$

C. $\left(1, \frac{5-\sqrt{5}}{2}\right)$

D. $\left[1, \frac{5+\sqrt{5}}{2}\right)$

Answer: C

Solution:

Solution:
$-1 \le 2x^2 - 3 < 2$
or $2 \le 2x^2 < 5$
or $1 \le x^2 < \frac{5}{2}$
$x \in \left(-\sqrt{\frac{5}{2}}, -1\right] \cup \left[1, \sqrt{\frac{5}{2}}\right)$
$\log_{\frac{1}{2}}(x^2 - 5x + 5) > 0$
$0 < x^2 - 5x + 5 < 1$
$x^2 - 5x + 5 > 0 \& x^2 - 5x + 4 < 0$
$x \in \left(-\infty, \frac{5-\sqrt{5}}{2}\right) \cup \left(\frac{5+\sqrt{5}}{2}, \infty\right)$
$\& x \in (-\infty, 1) \cup (4, \infty)$
Taking intersection
$x \in \left(1, \frac{5-\sqrt{5}}{2}\right)$

Question96

The number of functions f , from the set $A = \{x \in N : x^2 - 10x + 9 \le 0\}$ to the set $B = \{n^2 : n \in N\}$ such that $f(x) \le (x-3)^2 + 1$, for every $x \in A$, is _____. [27-Jul-2022-Shift-2]

Answer: 1440

Solution:

Solution:
$A = \{x \in N, x^2 - 10x + 9 \le 0\}$
$= \{1, 2, 3, ..., 9\}$
$B = \{1, 4, 9, 16, ...\}$

$f(x) \le (x-3)^2 + 1$
$f(1) \le 5, f(2) \le 2, \ldots\ldots\ldots f(9) \le 37$
$x = 1$ has 2 choices
$x = 2$ has 1 choice
$x = 3$ has 1 choice
$x = 4$ has 1 choice
$x = 5$ has 2 choices
$x = 6$ has 3 choices
$x = 7$ has 4 choices
$x = 8$ has 5 choices
$x = 9$ has 6 choices
$\therefore$ Total functions $= 2 \times 1 \times 1 \times 1 \times 2 \times 3 \times 4 \times 5 \times 6 = 1440$

Question97

Considering only the principal values of the inverse trigonometric functions, the domain of the function $f(x) = \cos^{-1}\left(\frac{x^2 - 4x + 2}{x^2 + 3}\right)$ is :

[28-Jul-2022-Shift-1]

Options:

A. $\left(-\infty, \frac{1}{4}\right]$

B. $\left[-\frac{1}{4}, \infty\right)$

C. $(-1/3, \infty)$

D. $\left(-\infty, \frac{1}{3}\right]$

Answer: B

Solution:

Solution:
$-1 \le \frac{x^2 - 4x + 2}{x^2 + 3} \le 1$
$\Rightarrow -x^2 - 3 \le x^2 - 4x + 2 \le x^2 + 3$
$\Rightarrow 2x^2 - 4x + 5 \ge 0 \quad -4x \le 1$

$x \in R \& x \geq -\frac{1}{4}$

So domain is $\left[-\frac{1}{4}, \infty\right)$

Question98

Let α, β and γ be three positive real numbers. Let $f(x) = \alpha x^5 + \beta x^3 + \gamma x$, $x \in R$ and $g : R \to R$. be such that $g(f(x)) = x$ for all $x \in R$. If $a_1, a_2, a_3, \ldots, a_n$ be in arithmetic progression with mean zero, then the value of $f\left(g\left(\frac{1}{n}\sum_{i=1}^{n} f(a_i)\right)\right)$ is equal to:

[28-Jul-2022-Shift-1]

Options:

A. 0

B. 3

C. 9

D. 27

Answer: A

Solution:

Solution:

$$f\left(g\left(\frac{1}{n}\sum_{i=1}^{n} f(a_i)\right)\right)$$

$$\frac{a_1 + a_2 + a_3 + \ldots\ldots + a_n}{n} = 0$$

∴ First and last term, second and second last and so on are equal in magnitude but opposite in sign.

$$f(x) = \alpha x^5 + \beta x^3 + \gamma x$$

$$\sum_{i=1}^{n} f(a_i) = \alpha(a_1^5 + a_2^5 + a_3^5 + \ldots + a_n^5) + \beta(a_1^3 + a_2^3 + \ldots + a_n^3) + \gamma(a_1 + a_2 + \ldots + a_n)$$

$$= 0\alpha + 0\beta + 0\gamma$$

$$= 0$$

$$\therefore f\left(g\left(\frac{1}{n}\sum_{i=1}^{n} f(a_i)\right)\right) = \frac{1}{n}\sum_{i=1}^{n} f(a_i) = 0$$

Question99

The number of elements in the set

$S = \left\{ x \in \mathbb{R} : 2\cos\left(\frac{x^2+x}{6} \right) = 4^x + 4^{-x} \right\}$ is:

[29-Jul-2022-Shift-2]

Options:

A. 1

B. 3

C. 0

D. infinite

Answer: A

Solution:

Solution:

$2\cos\left(\frac{x^2+x}{6} \right) = 4^x + 4^{-x}$

L.H.S ≤ 2. & R.H.S. ≥ 2

Hence L.H.S $= 2$& R.H.S $= 2$

$2\cos\left(\frac{x^2+x}{6} \right) = 2 \;\; 4^x + 4^{-x} = 2$

Check $x = 0$ Possible hence only one solution.

Question100

The domain of the function $f(x) = \sin^{-1}\left(\frac{x^2-3x+2}{x^2+2x+7} \right)$ is:

[29-Jul-2022-Shift-2]

Options:

A. $[1, \infty)$

B. $[-1, 2]$

C. $[-1, \infty)$

D. $(-\infty, 2]$

Answer: C

Solution:

Solution:

$$f(x) = \sin^{-1}\left(\frac{x^2-3x+2}{x^2+2x+7}\right)$$

$$-1 \le \frac{x^2-3x+2}{x^2+2x+7} \le 1$$

$$\frac{x^2-3x+2}{x^2+2x+7} \le 1$$

$$x^2-3x+2 \le x^2+2x+7$$

$$5x \ge -5$$

$$x \ge -1$$

And $\frac{x^2-3x+2}{x^2+2x+7} \ge -1$

$$x^2-3x+2 \ge -x^2-2x-7$$

$$2x^2-x+9 \ge 0$$

$x \in R$

(i) $\cap$ (ii)

Domain $\in [-1, \infty)$

Question101

The total number of functions, f : {1, 2, 3, 4} → {1, 2, 3, 4, 5, 6} such that f (1) + f (2) = f (3), is equal to : [25-Jul-2022-Shift-1]

Options:

A. 60

B. 90

C. 108

D. 126

Answer: B

Solution:

Solution:
Given, $f(1)+f(2)=f(3)$
It means $f(1)$, $f(2)$ and $f(3)$ are dependent on each other. But there is no condition on $f(4)$, so $f(4)$ can be $f(4)=1, 2, 3, 4, 5, 6$.
For $f(1)$, $f(2)$ and we have to find how many functions possible which will satisfy the condition $f(1)+f(2)=f(3)$
Case 1:
When $f(3)=2$ then possible values of $f(1)$ and $f(2)$ which satisfy $f(1)+f(2)=f(3)$ is $f(1)=1$ and $f(2)=1$.
And $f(4)$ can be $=1, 2, 3, 4, 5, 6$
∴ Total possible functions $=1\times 6=6$
Case 2 :

When $f(3)=3$ then possible values
(1) $f(1)=1$ and $f(2)=2$
(2) $f(1)=2$ and $f(2)=1$
And $f(4)$ can be $=1, 2, 3, 4, 5, 6$.
∴ Total functions $=2\times 6=12$
Case 3 :
When $f(3)=4$ then
(1) $f(1)=1$ and $f(2)=3$
(2) $f(1)=2$ and $f(2)=2$
(3) $f(1)=3$ and $f(2)=1$
And $f(4)$ can be $=1, 2, 3, 4, 5, 6$
∴ Total functions $=3\times 6=18$
Case 4 :
When $f(3)=5$ then
(1) $f(1)=1$ and $f(4)=4$
(2) $f(1)=2$ and $f(4)=3$
(3) $f(1)=3$ and $f(4)=2$
(4) $f(1)=4$ and $f(4)=1$
And $f(4)$ can be $=1, 2, 3, 4, 5$ and 6
∴ Total functions $=4\times 6=24$
Case 5 :
When $f(3)=6$ then
(1) $f(1)=1$ and $f(2)=5$
(2) $f(1)=2$ and $f(2)=4$
(3) $f(1)=3$ and $f(2)=3$
(4) $f(1)=4$ and $f(2)=2$
(5) $f(1)=5$ and $f(2)=1$
And $f(4)$ can be $=1, 2, 3, 4, 5$ and 6
∴ Total possible functions $=5\times 6=30$
∴ Total functions from those 5 cases we get $=6+12+18+24+30=90$

Question102

Let f : N → R be a function such that f (x + y) = 2f (x)f (y) for natural numbers x and y. If f(1) = 2, then the value of α for which

$\sum_{k=1}^{10} f(\alpha + k) = \frac{512}{3}(2^{20} - 1)$
holds, is :
[25-Jun-2022-Shift-1]

Options:

A. 2

B. 3

C. 4

D. 6

Answer: C

Solution:

Solution:

Given,

$f(x+y) = 2f(x)f(y)$

and $f(1) = 2$

For $x = 1$ and $y = 1$,

$f(1+1) = 2f(1)f(1)$

$\Rightarrow f(2) = 2(f(1))^2 = 2(2)^2 = 2^3$

For $x = 1, y = 2$

$f(1+2) = 2f(1)y(2)$

$\Rightarrow f(3) = 2 \cdot 2 \cdot 2^3 = 2^5$

For $x = 1, y = 3$

$f(1+3) = 2f(1)f(3)$

$\Rightarrow f(4) = 2 \cdot 2 \cdot 2^5 = 2^7$

For $x = 1, y = 4$

$f(1+4) = 2f(1)f(4)$

$\Rightarrow f(5) = 2.2 \cdot 2^7 = 2^9 \ldots.$

Also given

$$\sum_{k=1}^{10} f(\alpha + k) = \frac{512}{3}(2^{20} - 1)$$

$$\Rightarrow f(\alpha+1) + f(\alpha+2) + f(\alpha+3) + \ldots + f(\alpha+10) = \frac{512}{3}(2^{20} - 1)$$

$$\Rightarrow f(\alpha+1) + f(\alpha+2) + f(\alpha+3) + \ldots + f(\alpha+10) = \frac{2^9((2^2)^{10} - 1)}{2^2 - 1}$$

This represent a G.P with first term $= 2^9$ and common ratio $= 2^2$

$\therefore$ First term $= f(\alpha+1) = 2^9 \ldots . (2)$

From equation $(1), f(5) = 2^9$

$\therefore$ From (1) and (2), we get

$f(\alpha+1) = 2^9 = f(5)$

$\Rightarrow f(\alpha+1) = f(5)$

$\Rightarrow f(\alpha+1) = f(4+1)$

Comparing both sides we get, $\alpha = 4$

Question103

Let f : R → R be a function defined by

$f(x) = \left(2\left(1 - \frac{x^{25}}{2}\right)(2 + x^{25})\right)^{\frac{1}{50}}$. If the function

$g(x) = f(f(f(x)) + f(f(x))$, then the greatest integer less than or equal to g(1) is___
[25-Jun-2022-Shift-1]

Answer: 2

Solution:

Solution:

Given,

$$f(x) = \left(2\left(1 - \frac{x^{25}}{2}\right)(2 + x^{25})\right)^{\frac{1}{50}}$$

and $g(x) = f(f(f(x))) + f(f(x))$

$\therefore g(1) = f(f(f(1))) + f(f(1))$

Now, $f(1) = \left(2\left(1 - \frac{1^{25}}{2}\right)(2 + 1^{25})\right)^{\frac{1}{50}}$

$$= \left(2\left(1 - \frac{1}{2}\right)(2 + 1)\right)^{\frac{1}{50}}$$

$$= (3)^{\frac{1}{50}}$$

$\therefore f(f(1)) = f\left(3^{\frac{1}{50}}\right)$

$$= \left(2\left(1 - \frac{\left(3^{\frac{1}{50}}\right)^{25}}{2}\right)\left(2 + \left(3^{\frac{1}{50}}\right)^{25}\right)\right)^{\frac{1}{50}}$$

$$= \left(2\left(1 - \frac{3^{\frac{1}{2}}}{2} \right)\left(2 + 3^{\frac{1}{2}} \right) \right)^{\frac{1}{50}}$$

$$= \left(2 \times \left(\frac{2 - \sqrt{3}}{2} \right)(2 + \sqrt{3}) \right)^{\frac{1}{50}}$$

$$= [(2 - \sqrt{3})(2 + \sqrt{3})]^{\frac{1}{50}}$$

$$= (4 - 3)^{\frac{1}{50}}$$

$$= 1^{\frac{1}{50}} = 1$$

Now, $f(f(f(1))) = f(1) = 3^{\frac{1}{50}}$

$\therefore g(1) = f(f(f(1))) + f(f(1))$

$$= 3^{\frac{1}{50}} + 1$$

Now, greatest integer less than or equal to $g(1)$

$$= [g(1)]$$

$$= \left[3^{\frac{1}{50}} + 1 \right]$$

$$= \left[3^{\frac{1}{50}} \right] + [1]$$

$$= [1.02] + 1$$

$$= 1 + 1 = 2$$

Question104

Let $f(x) = \frac{x-1}{x+1}$, $x \in R - \{0, -1, 1\}$. If $f^{n+1}(x) = f(f^n(x))$ for all $n \in N$, then $f^6(6) + f^7(7)$ is equal to :
[26-Jun-2022-Shift-1]

Options:

A. $\frac{7}{6}$

B. $-\frac{3}{2}$

C. $\frac{7}{12}$

D. $-\frac{11}{12}$

Answer: B

Solution:

Solution:

Given,

$$f(x) = \frac{x-1}{x+1}$$

Also given,

$$f^{n+1}(x) = f(f^n(x)) \ldots . (1)$$

$\therefore$ For $n = 1$

$$f^{1+1}(x) = f(f^1(x))$$

$$\Rightarrow f^2(x) = f(f(x))$$

$$= f\left(\frac{x-1}{x+1}\right)$$

$$= \frac{\frac{x-1}{x+1}-1}{\frac{x-1}{x+1}+1}$$

$$= \frac{\frac{x-1-x-1}{x+1}}{\frac{x-1+x+1}{x+1}}$$

$$= \frac{-2}{2x} = -\frac{1}{x}$$

From equation (1), when $n = 2\ f^{2+1}(x) = f(f^2(x))$

$$\Rightarrow f^3(x) = f(f^2(x))$$

$$= f\left(-\frac{1}{x}\right)$$

$$= \frac{-\frac{1}{x}-1}{-\frac{1}{x}+1}$$

$$= \frac{\frac{-1-x}{-1}}{\frac{-1+x}{x}}$$

$$= \frac{-1-x}{-1+x} = \frac{-(x+1)}{x-1}$$

Similarly,

$$f^4(x) = f(f^3(x))$$

$$= f\left(\frac{-x+1}{x-1}\right)$$

$$\frac{-(x+1)}{} - 1$$

$$= \frac{x-1}{\frac{-(x+1)}{x-1}+1}$$

$$= \frac{\frac{x-1-x+1}{x-1}}{\frac{-x-1+x-1}{x-1}}$$

$$= \frac{-2x}{-2} = x$$

$$\therefore f^5(x) = f(f^4(x))$$

$$= f(x)$$

$$= \frac{x-1}{x+1}$$

$$f^6(x) = f(f^5(x))$$

$$= f\left(\frac{x-1}{x+1}\right)$$

$$= -\frac{1}{x}$$ (Already calculated earlier)

$$f^7(x) = f(f^6(x))$$

$$= f\left(-\frac{1}{x}\right)$$

$$-\frac{1}{x} - 1$$

$$= \frac{x}{-\frac{1}{x}+1}$$

$$= \frac{-(x+1)}{x-1}$$

$$\therefore f^6(6) = -\frac{1}{6}$$

and $f^7(7) = \frac{-(7+1)}{7-1} = -\frac{8}{6}$

So, $f^6(6) + f^7(7)$

$$= -\frac{1}{6} - \frac{8}{6}$$

$$= -\frac{3}{2}$$

Question105

The range of the function,

$$\mathbf{f(x) = \log_{\sqrt{5}}\left(3 + \cos\left(\frac{3\pi}{4} + x\right) + \cos\left(\frac{\pi}{4} + x\right) + \cos\left(\frac{\pi}{4} - x\right) - \cos\left(\frac{3\pi}{4} - x\right)\right)}$$

is

[2021, 01 Sep. Shift-II]

Options:

A. $(0, \sqrt{5})$

B. $[-2, 2]$

C. $\left[\frac{1}{\sqrt{5}}, \sqrt{5}\right]$

D. [0, 2]

Answer: D

Solution:

Solution:

$$f(x)=\log_{\sqrt{5}}\left(3+\cos\left(\frac{3\pi}{4}+x\right)+\cos\left(\frac{\pi}{4}+x\right)\right.$$
$$\left.+\cos\left(\frac{\pi}{4}-x\right)-\cos\left(\frac{3\pi}{4}-x\right)\right)$$
$$=\log_{\sqrt{5}}(3-\sqrt{2}\sin x+\sqrt{2}\cos x)$$
$$\because -2\le -\sqrt{2}\sin x+\sqrt{2}\cos x\le 2$$
$$\Rightarrow 1\le 3-\sqrt{2}\sin x+\sqrt{2}\cos x\le 5$$
$$\Rightarrow \log_{\sqrt{5}}1\le \log_{\sqrt{5}}(3-\sqrt{2}\sin x+\sqrt{2}\cos x)$$
$$\Rightarrow 0\le f(x)\le 2$$
$$\Rightarrow f(x)\in[0,2]$$

Question106

The domain of the function $f(x)=\sin^{-1}\left(\frac{3x^2+x-1}{(x-1)^2}\right)+\cos^{-1}\left(\frac{x-1}{x+1}\right)$ is

[2021, 31 Aug. Shift-II]

Options:

A. $\left[0,\ \frac{1}{4}\right]$

B. $[-2,0]\cup\left[\frac{1}{4},\ \frac{1}{2}\right]$

C. $\left[\frac{1}{4},\ \frac{1}{2}\right]\cup\{0\}$

D. $\left[0,\ \frac{1}{2}\right]$

Answer: C

Solution:

Solution:

$$f(x)=\sin^{-1}\left(\frac{3x^2+x-1}{(x-1)^2}\right)+\cos^{-1}\left(\frac{x-1}{x+1}\right)$$
$$-1\le\frac{x-1}{x+1}\le 1\Rightarrow -1-1\le\frac{x-1}{x+1}-1\le 1-1$$
$$\Rightarrow\ -2\le\frac{-2}{x+1}\le 0\Rightarrow 0\le\frac{1}{x+1}\le 1$$

$\Rightarrow\ 1 \le x+1 < \infty$
$\Rightarrow\ 0 \le x < \infty$
$\Rightarrow\ x \in [0, \infty)$
and $-1 \le \dfrac{3x^2+x-1}{(x-1)^2} \le 1$
$\Rightarrow\ -(x-1)^2 \le 3x^2+x-1 \le (x-1)^2,\ x \ne 1$
$\Rightarrow\ -(x^2-2x+1) \le 3x^2+x-1$
and $3x^2+x-1 \le x^2-2x+1$
$\Rightarrow\ 4x^2-x \ge 0$
and $2x^2+3x-2 \le 0$
$\Rightarrow\ x(4x-1) \ge 0$
and $(x+2)(2x-1) \le 0$
$\Rightarrow\ x \in (-\infty, 0] \cup \left[\dfrac{1}{4}, \infty\right)$
and $x \in \left[-2, \dfrac{1}{2}\right]$
$\Rightarrow\ x \in (-2, 0] \cup \left[\dfrac{1}{4}, \dfrac{1}{2}\right]$
Domain of f in Eq. (i) $\cap$ Eq. (ii)
$\therefore\ x \in \{0\} \cup \left[\dfrac{1}{4}, \dfrac{1}{2}\right]$

Question107

Let $f : N \to N$ be a function such that $f(m+n) = f(m) + f(n)$ for every $m, n \in N$. If $f(6) = 18$, then $f(2) \cdot f(3)$ is equal to [2021, 31 Aug. Shift-11]

Options:

A. 6

B. 54

C. 18

D. 36

Answer: B

Solution:

Solution:
$f(m+n) = f(m) + f(n),\ m, n \in N$
$\therefore\ f(3+3) = f(3) + f(3)$

$\Rightarrow \quad f(6) = 2f(3) = 18 \quad [\because f(6) = 18]$
Also $f(3) = f(2+1) = f(2) + f(1)$
$= f(1+1) + f(1)$
$f(3) = f(1) + f(1) + f(1)$
$\Rightarrow \quad 9 = 3f(1) \Rightarrow f(1) = 3$
$\therefore \quad f(2) = f(1+1) = f(1) + f(1) = 3 + 3 = 6$
Hence, $f(2) \cdot f(3) = 6 \cdot 9 = 54$

Question108

The domain of the function $\text{cosec}^{-1}\left(\frac{1+x}{x}\right)$ is

[2021, 26 Aug. Shift-II]

Options:

A. $\left(-1, -\frac{1}{2}\right] \cup (0, \infty)$

B. $\left[-\frac{1}{2}, 0\right) \cup [1, \infty)$

C. $\left(-\frac{1}{2}, \infty\right) - \{0\}$

D. $\left[-\frac{1}{2}, \infty\right) - \{0\}$

Answer: D

Solution:

Solution:
$f(x) = \text{cosec}^{-1}\left(\frac{1+x}{x}\right) \quad \left|\frac{1+x}{x}\right| \geq 1$
Clearly, $x \neq 0$
$r\,|1+x|^2 \geq |x|^2$
$1 + x^2 + 2x \geq x^2$
$2x + 1 \geq 0$
$x \geq -\frac{1}{2}$
So,
$x \in \left[-\frac{1}{2}, \infty\right] - \{0\}$

Question109

Which of the following is not correct for relation R on the set of real numbers ?
[2021, 31 Aug. Shift-1]

Options:

A. $(x, y) \in R \Leftrightarrow 0 < |x| - |y| \leq 1$ is neither transitive nor symmetric.

B. $(x, y) \in R \Leftrightarrow 0 < |x - y| \leq 1$ is symmetric and transitive.

C. $(x, y) \in R \Leftrightarrow x| - |y| \leq 1$ is reflexive but not symmetric.

D. $(x, y) \in R \Leftrightarrow x - y| \leq 1$ is reflexive and symmetric.

Answer: B

Solution:

Solution:
According to the question, let's consider option (b) $(2, 3)$ and $(3, 4)$ satisfy $0 < |x - y| \text{ leq } 1$ but $(2, 4)$ does not satisfy it.

Question110

Let N be the set of natural numbers and a relation R on N be defined by $R = \{(x, y) \in N \times N : x^3 - 3x^2y - xy^2 + 3y^3 = 0\}$.Then the relation R is
[2021, 27 July Shift-11]

Options:

A. symmetric but neither reflexive nor transitive.

B. reflexive but neither symmetric nor transitive.

C. reflexive and symmetric, but not transitive.

D. an equivalence relation.

Answer: B

Solution:

Solution:
Given, relation R on N is defined by
$R = \{(x, y) \in N \times N : x^3 - 3x^2 - xy^2 + 3y^3 = 0\}$
$x^3 - 3x^2y - xy^2 + 3y^3 = 0$
$\Rightarrow x^3 - xy^2 - 3x^2y + 3y^3 = 0$
$\Rightarrow x(x^2 - y^2) - 3y(x^2 - y^2) = 0$
$\Rightarrow (x - 3y)(x^2 - y^2) = 0$
$\Rightarrow (x - 3y)(x - y)(x + y) = 0$
Now, $x - x = 0$
$\Rightarrow x = x, \forall (x, x) \in N \times N$
So, R is a reflexive relation.
But not symmetric and transitive relation because,
$(3, 1)$ satisfies but $(1, 3)$ does not. Also, $(3, 1)$ and
$(1, -1)$ satisfies but $(3, -1)$ does not.
Hence, relation R is reflexive but neither symmetric nor transitive.

Question111

Define a relation R over a class of n × n real matrices A and B as "ARB, if there exists a non-singular matrix P such that $PAP^{-1} = B'$. Then which of the following is true ? [2021, 18 March Shift-II]

Options:

A. R is symmetric, transitive but not reflexive.

B. R is reflexive, symmetric but not transitive.

C. R is an equivalence relation.

D. R is reflexive, transitive but not symmetric.

Answer: C

Solution:

Solution:

For reflexive relation, $\forall (A, A) \in R$ for matrix P.
$\Rightarrow A = PAP^{-1}$ is true for $P = 1$
So, R is reflexive relation.
For symmetric relation,
Let $(A, B) \in R$ for matrix P.
$\Rightarrow \ A = PBP^{-1}$After pre-multiply by P^{-1} and post-multiply by P_1
we get
$P^{-1}AP = B$
So, $(B, A) \in R$ for matrix P^{-1}.
So, R is a symmetric relation.
For transitive relation,
Let ARB and BRC
So, $A = PBP^{-1}$ and $B = PCP^{-1}$
Now, $\ A = P(PCP^{-1})P^{-1}$
$\Rightarrow \ A = (P)^2C(P^{-1})^2 \Rightarrow A = (P)^2 \cdot C \cdot (P^2)^{-1}$
$\therefore (A, C) \in R$ for matrix P^2.
$\therefore R$ is transitive relation.
Hence, R is an equivalence relation.

Question112

Let A = {2, 3, 4, 5, …., 30} and ' $''$ be an equivalence relation on A × A, defined by (a, b) ~ (c, d), if and only if ad = bc. Then, the number of ordered pairs, which satisfy this equivalence relation with ordered pair (4, 3) is equal to
[2021, 16 March Shift-II]

Options:

A. 5

B. 6

C. 8

D. 7

Answer: D

Solution:

Solution:
$A = \{2, 3, 4, 5, \ldots, 30\}$
$a = bc$

$\therefore$ (a, b)R(4, 3)
$\Rightarrow$ 3a = 4b
$\Rightarrow$ $a = \left(\frac{4}{3}\right) b$
b must be a multiple of 3, b can be
(3, 6, 9, .. . 30).
Also, a must be less than or equal to 30.
(a, b) = (4, 3), (8, 6), (12, 9), (16, 12), (20, 15)
(24, 18), (28, 21)
$\Rightarrow$7 ordered pairs

Question113

Let R = {(P, O)|, P and Q are at the same distance from the origin\} be a relation, then the equivalence class of (1, −1) is the set [2021, 26 Feb. Shift-1]

Options:

A. $S = \{(x, y) \mid x^2 + y^2 = 4\}$

B. $S = \{(x, y) \mid x^2 + y^2 = 1\}$

C. $S = \{(x, y) \mid x^2 + y^2 = \sqrt{2}\}$

D. $S = \{(x, y) \mid x^2 + y^2 = 2\}$

Answer: D

Solution:

Solution:
Let P(a, b) and Q(c, d) are any two points.
Given, OP = 00
i.e. $\sqrt{a^2 + b^2} = \sqrt{c^2 + d^2}$
Squaring on both sides,
$R = \{((a, b), (c, d)) : a^2 + b^2 = c^2 + d^2\}$
R(x, y), S(1, −1), OR = OS (equivalence class)
This gives OR = $\sqrt{x^2 + y^2}$ and OS = $\sqrt{2}$
1 $\Rightarrow$ $\sqrt{x^2 + y^2} = \sqrt{2}$
$\Rightarrow x^2 + y^2 = 2$(Squaring on both sides)
$\therefore$ $S = \{(x, y) : x^2 + y^2 = 2\}$

Question114

Let {S = 1, 2, 3, 4, 5, 6, 7}. Then the number of possible functions $f : S \to S$ such that $f(m \cdot n) = f(m) \cdot f(n)$ for every $m, n \in S$ and $m \cdot n \in S$ is equal to .....
[2021, 27 July Shift-I]

Answer: 490

Solution:

Solution:

$S = \{1, 2, 3, 4, 5, 6, 7\}$

$f : S \to S$

$f(m \cdot n) = f(m)f(n)$

$m, n \in S \Rightarrow m, n \in S$

If $mn \in S \Rightarrow mn \leq 7$

So, $(1 \cdot 1, 1 \cdot 2, \ldots, 1 \cdot 7) \leq 7$

$(2 \cdot 2, 2 \cdot 3) \leq 7$

When $m = 1$, $f(n) = f(1) \cdot f(n) \Rightarrow f(1) = 1$

When $m = n = 2$,

$$f(4) = f(2)f(2) = \begin{cases} f(2) = 1 \Rightarrow f(4) = 1 \text{ or} \\ f(2) = 2 \Rightarrow f(4) = 4. \end{cases}$$

When, $m = 2, n = 3$

$$f(6) = f(2)f(3) \begin{cases} \text{When, } f(2) = 1 \\ f(3) = 1 \text{ to } 7 \\ \text{When, } f(2) = 2 \\ f(3) = 1 \text{ or } 2 \text{ or } 3. \end{cases}$$

And $f(5), f(7)$ can take any value (1-7) [$\because f(5) = f(1) \cdot f(5) \leq 7$ and $f(7) = f(1) \cdot f(7) \leq 7$] The possible combination is

ll $f(1) = 1$ $f(1) = 1$

$f(2) = 1$ $f(2) = 2$

$f(3) = (1 - 7)$ $f(3) = (1 - 3)$

$f(4) = 1$ $f(4) = 4$

$f(5) = (1 - 7)$ $f(5) = (1 - 7)$

$f(6) = f(3)$ $f(6) = f(3)$

$f(7) = (1 - 7)$ $f(7) = (1 - 7)$

So, total $= (1 \times 1 \times 7 \times 1 \times 7 \times 1 \times 7)$

$+(1 \times 1 \times 3 \times 1 \times 7 \times 1 \times 7)$

$= 490$

Question115

If [x] be the greatest integer less than or equal to x, then $\sum_{n=8}^{100}\left[\frac{(-1)^n n}{2}\right]$ is equal to
[25 July 2021, Shift-III]

Options:

A. 0

B. 4

C. -2

D. 2

Answer: D

Solution:

Solution:
We have,
$\sum_{n=8}^{100}\left[\frac{(-1)^n n}{2}\right]$ ($\because$[x] is the greatest integer function)
Substitute the values of n

$$= [4]+[-4.5]+[5]+[-5.5] + \ldots +[-49.5]+[50]$$
$$= 4-5+5-6+\ldots-50+50$$
$$= 4$$

Question116

If the domain of the function f (x) = $\frac{\cos^{-1}\sqrt{x^2-x+1}}{\sqrt{\sin^{-1}\left(\frac{2x-1}{2}\right)}}$ is the interval

(α, β), then α + β is equal to
[2021, 22 July Shift-II]

Options:

A. $\frac{3}{2}$

B. 2

C. $\frac{1}{2}$

D. 1

Answer: A

Solution:

Solution:

$$f(x) = \frac{\cos^{-1}\sqrt{x^2 - x + 1}}{\sqrt{\sin^{-1}\left(\frac{2x-1}{2}\right)}}$$

$\Rightarrow x \in R, \quad x(x-1) \leq 0$

$x^2 - x + 1 \geq 0$ and $x^2 - x + 1 \leq 1$

$0 \leq x \leq 1 \quad \cdots\cdots (i) \Rightarrow \quad 0 < \sin^{-1}\left(\frac{2x-1}{2}\right) < \frac{\pi}{2}$

$\Rightarrow \quad 0 < \frac{2x-1}{2} < 1$

$\Rightarrow \quad \frac{1}{2} < x < \frac{3}{2} \quad \cdots\cdots (ii)$

$(A) \cap (B) \quad = x \in \left(\frac{1}{2}, 1\right]$

$\therefore \quad \alpha + \beta \quad = \frac{3}{2}$

Question117

Let [x] denote the greatest integer ≤x, where x ∈ R. If the domain of the real valued function

$f(x) = \sqrt{\frac{|[x]| - 2}{|[x]| - 3}}$ is $(-\infty, a) \cup [b, c) \cup [u, \infty)$, a < b < c, then the value of a + b + c is

[2021, 20 July Shift-I]

Options:

A. 8

B. 1

C. −2

D. −3

Answer: C

Solution:

Solution:

$f(x) = \sqrt{\frac{|[x]|-2}{|[x]|-3}\,\frac{|[x]|-2}{|[x]|-3}} \geq 0$

Let $|[x]| = t$



1 $|[x]| = 3 \Rightarrow x \in [-3, -2) \cup [3, 4)$

Domain of $x = [-\infty, -3) \cup [-2, 3) \cup [4, \infty)$

$a = -3$

$b = -2$

$c = 3$

$\therefore\ a + b + c = -3 + (-2) + 3 = -2$

Question118

The real valued function $f(x) = \frac{\text{cosec}^{-1}x}{\sqrt{x - [x]}}$, where [x] denotes the greatest integer less than or equal to x, is defined for all x belonging to [2021, 18 March Shift-I]

Options:

A. all reals except integers

B. all non-integers except the interval $[-1, 1]$

C. all integers except $0, -1, 1$

D. all reals except the interval $[-1, 1]$

Answer: B

Solution:

Solution:

Given, $f(x) = \frac{\text{cosec}^{-1}x}{\sqrt{x-[x]}}$

$\Rightarrow f(x) = \frac{\text{cosec}^{-1}x}{\sqrt{\{x\}}}$

For $f(x)$ to be defined,

$$\begin{cases} |x| \geq 1 \\ \{x\} > 0. \end{cases} \Rightarrow \begin{cases} x \leq -1 \text{ or } x \geq 1 \\ x \neq 1 \text{ integers } . \end{cases}$$

i.e. $x \in (-\infty, -1] \cup [1, \infty) - \{$ integers $\}$

i.e. all non-integers except the interval $[-1, 1]$

(here, -1 and 1 are included in except case, because of -1 and 1 are integers).

Question119

If the functions are defined as $f(x) = \sqrt{x}$ and $g(x) = \sqrt{1-x}$, then what is the common domain of the following functions?

$f + g, f - g, f / g, g / f, g - f$, where

$(f \pm g)(x) = f(x) \pm g(x), (f / g)(x) = \frac{f(x)}{g(x)}$

[2021, 18 March, Shift-1]

Options:

A. $0 \leq x \leq 1$

B. $0 \leq x < 1$

C. $0 < x < 1$

D. $0 < x \leq 1$

Answer: C

Solution:

Solution:
Given, $f(x) = \sqrt{x}$ and $g(x) = \sqrt{1-x}$
$\therefore$ Domain of $f(x) = D_1$ is $x \geq 0$
i.e. $D_1 : x \in (0, \infty)$
and domain of $g(x) = D_2$ is $1 - x \geq 0 \Rightarrow x \leq 1$
i.e. $D_2 : x \in (-\infty 1]$
As, we know that, the domain of $f + g_1 f - g, g - f$ will be $D_1 \cap D_2$ as well as the domain for $\frac{f}{g}$ is $D_1 \cap D_2$ except all
those value(s) of x, such that $g(x) = 0$.
Similarly, for $\frac{g}{f}$ is $D_1 \cap D_2$ but $f(x) \neq 0$.
Hence, common domain for $(f + g), (f - g), \left(\frac{f}{g}\right), \left(\frac{g}{f}\right)$ and $(g - f)$ will be $0 < x < 1$

Question120

A function $f(x)$ is given by $f(x) = \frac{5^x}{5^x + 5}$, then the sum of the series $f\left(\frac{1}{20}\right) + f\left(\frac{2}{20}\right) + f\left(\frac{3}{20}\right) + \ldots + f\left(\frac{39}{20}\right)$ is equal to [2021, 25 Feb. Shift-II]

Options:

A. $\frac{29}{2}$

B. $\frac{49}{2}$

C. $\frac{39}{2}$

D. $\frac{19}{2}$

Answer: C

Solution:

Solution:

Given, $f(x)=\frac{5^x}{5^x+5}$, then,

$f(2-x)=\frac{5^{2-x}}{5^{2-x}+5}$

$=\frac{5}{5^x+5}$

This gives, $f(x)+f(2-x)=\frac{5^x+5}{5^x+5}=1 \Rightarrow f\left(\frac{1}{20}\right)+f\left(2-\frac{1}{20}\right)=f\left(\frac{1}{20}\right)+f\left(\frac{39}{20}\right)=1$

Similarly,

$cf\left(\frac{2}{20}\right)+f\left(\frac{38}{20}\right)=1$ and so on,

$\therefore f\left(\frac{1}{20}\right)+f\left(\frac{2}{20}\right)+\ldots+f\left(\frac{38}{20}\right)+f\left(\frac{39}{20}\right)$

$=1+1+\ldots+1+f\left(\frac{20}{20}\right)$

$=19+f(1)=19+\frac{1}{2}=\frac{39}{2}$

Question121

If $a+\alpha=1$, $b+\beta=2$ and $af(x)+\alpha f\left(\frac{1}{x}\right)=bx+\frac{\beta}{x}$, $x\neq 0$, then the value of expression $\frac{f(x)+f\left(\frac{1}{x}\right)}{x+\frac{1}{x}}$ is

[2021, 24 Feb. Shift-II]

Answer: 2

Solution:

Solution:

Given, $a+\alpha=1$

$b+\beta=2$

$\because a\cdot f(x)+\alpha\cdot f\left(\frac{1}{x}\right)=bx+\frac{\beta}{x} \quad \cdots\cdots\cdot(i)$

Replace x by $\frac{1}{x}$,

$af\left(\frac{1}{x}\right)+af(x)=\frac{b}{x}+\beta x$

Adding Eqs. (i) and (ii), we get

$$\&(a+\alpha)\left[f(x)+f\left(\frac{1}{x}\right)\right]=\left(x+\frac{1}{x}\right)(b+\beta) \quad \cdots\cdots\cdot (ii)$$

$$\Rightarrow \quad \frac{f(x)+f\left(\frac{1}{x}\right)}{x+\frac{1}{x}}=\frac{b+\beta}{a+\alpha}=\frac{2}{1}=2$$

Question122

Let $f(x) = \sin^{-1}x$ and $g(x) = \frac{x^2-x-2}{2x^2-x-6}$ If $g(2) = \lim_{x\to 2} g(x)$, then the domain of thefunction fog is [2021, 26 Feb. Shift-II]

Options:

A. $(-\infty, -2] \cup \left[-\frac{3}{2}, \infty\right)$

B. $(-\infty, -2] \cup [-1, \infty)$

C. $(-\infty, -2] \cup \left[-\frac{4}{3}, \infty\right)$

D. $(-\infty, -1] \cup [2, \infty)$

Answer: C

Solution:

Solution:

Given, $g(x) = \frac{x^2-x-2}{2x^2-x-6}$, $f(x) = \sin^{-1}x$

$$f(g(x)) = \sin^{-1}(g(x))$$

$$f \circ g(x) = \sin^{-1}\left(\frac{x^2-x-2}{2x^2-x-6}\right)$$

For the domain of $f \circ g(x)$,
$|g(x)| \leq 1$
[$\because$ Domain of $f(x)$ is $[-1, 1]$

$\Rightarrow \left| \frac{x^2 - x - 2}{2x^2 - x - 6} \right| \leq 1$

$\Rightarrow \left| \frac{(x+1)(x-2)}{(2x+3)(x-2)} \right| \leq 1$

$\Rightarrow \left| \frac{x+1}{2x+3} \right| \leq 1$

$\Rightarrow -1 \leq \frac{x+1}{2x+3} \leq 1$

$\Rightarrow \left(\frac{x+1}{2x+3} \right)^2 \leq 1$

$\Rightarrow (x+1)^2 \leq (2x+3)^2$

$\Rightarrow 3x^2 + 10x + 8 \geq 0$

$\Rightarrow (3x + y)(x + 2) \geq 0$

This implies,

$x \in (-\infty, -2] \cup \left[-\frac{4}{3}, \infty \right)$

Question123

Let g : N $\rightarrow$ N be defined as g(3n + 1) = 3n + 2 g(3n + 2) = 3n + 3, g(3n + 3) = 3n + 1, for all n $\geq$ 0. Then which of the following statements is true? [2021, 25 July Shift-1]

Options:

A. There exists an onto function f : N $\rightarrow$ N such that fog = f

B. There exists a one-one function f : N $\rightarrow$ N such that f$\circ$g = f

C. gogog = g

D. There exists a function f : N $\rightarrow$ N such that gof = f

Answer: A

Solution:

Solution:

$g(3n+1) = 3n+2$
$g(3n+2) = 3n+3$
$g(3n+3) = 3n+1$, for all $n \geq 0$
$g : N \rightarrow N$
$g(1) = 2, g(4) = 5, g(7) = 8$
$g(2) = 3, g(5) = 6, g(8) = 9$
$g(3) = 1, g(6) = 4, g(9) = 7$
$\Rightarrow \; f[g(1)] = f(1)$
$\Rightarrow \; f(2) = f(1)$
Clearly, it is not a one - one function.
Now, $f[g(2)] = f(2)$
$f(3) = f(2)$
And, $f[g(3)] = f(3)$
$f(1) = f(3)$
Similarly, $f[g(4)] = f(4)$
$f(5) = f(4)$
And, so on
$l\, f(1) = f(2) = f(3)$
$f(4) = f(5) = f(6)$



Now, there can be a possibility such that
So, $f(x)$ can be onto function.
When $f(1) = f(2) = f(3) = 1$
$f(4) = f(5) = f(6) = 2$
and so on.

Question124

Consider function $f : A \rightarrow B$ and $g : B \longrightarrow C(A, B, C \subset eqR)$ such that $(\ gof\ )^{-1}$ exists, then
[2021, 25 July Shift-II]

Options:

A. f and g both are one-one

B. f and g both are onto

C. f is one-one and g is onto

D. f is onto and g is one-one

Answer: C

Solution:

Solution:
Given functions, $f : A \to B$ and $g : B \to C(A, B, C \subset eqR)$
$\therefore (gof)^{-1}$ exists $\Rightarrow$ gof is a bijective function.
$\Rightarrow$ ' f ' must be 'one-one' and ' g ' must be 'onto' function.

Question125

Let A = {0, 1, 2, 3, 4, 5, 6, 7}. Then, the number of bijective functions F : A → A such that f (1) + f (2) = 3 − f (3) is equal to ..... [2021, 22 July Shift-III]

Answer: 720

Solution:

Solution:
$f(1) + f(2) = 3 - f(3)$
$A = \{0, 1, 2, 3, 4, 5, 6, 7\}$
$f : A \to A$
So, $f(1) + f(2) + f(3) = 3$
$0 + 1 + 2 = 3$ is the only possibility.
So, $f(0)$ can be either 0 or 1 or 2 .
Similarly, $f(1)$ and $f(2)$ can be 0,1 and 2 .
and $\{3, 4, 5, 6, 7\} \longrightarrow \{3, 4, 5, 6, 7\}$

They have 5 ! choices.
And $\{0, 1, 2\}$

They have 3! choices.
Number of bijective functions
$= 3! \times 5! = 720$

Question126

Let $f : R - \left\{ \frac{\alpha}{6} \right\} \rightarrow R$ be defined by

$f(x) = \frac{5x+3}{6x-\alpha}$

Then, the value of α for which (fof) (x) = x, for all $x \in R - \left\{ \frac{\alpha}{6} \right\}$ is

[2021, 20 July Shift-II]

Options:

A. No such α exists

B. 5

C. 8

D. 6

Answer: B

Solution:

Solution:

$f(x) = \frac{5x+3}{6x-\alpha}$

Now, $f \circ f(x) = f\left(\frac{5x+3}{6x-\alpha} \right)$

$= \frac{5\left(\frac{5x+3}{6x-\alpha} \right) + 3}{6\left(\frac{5x+3}{6x-\alpha} \right) - \alpha}$

$= \frac{5(5x+3)+3(6x-\alpha)}{6(5x+3)-\alpha(6x-2)}$

$= \frac{5(5x+3)+3(6x-\alpha)}{6(5x+3)-\alpha(6x-2)}$

Given, $fof(x) = x$

$\Rightarrow \frac{5(5x+3)+3(6x-\alpha)}{6(5x+3)-\alpha(6x-\alpha)} = x$

$\Rightarrow 25x + 15 + 18x - 3\alpha$

$= 30x^2 + 18x - 6\alpha x^2 + \alpha^2 x$

$\Rightarrow x^2(30-6\alpha) - x(\alpha^2 - 25) + 3\alpha - 15 = 0$

Comparing coefficients,

$30 - 6x = 0$

$\Rightarrow 6\alpha = 30$

$\Rightarrow \alpha = 5$

Question127

Let $f : R - \{3\} \to R - \{1\}$ be defined by $f(x) = \frac{x-2}{x-3}$. Let $g : R \to R$ be given as $g(x) = 2x - 3$. Then, the sum of all the values of x for which $f^{-1}(x) + g^{-1}(x) = \frac{13}{2}$ is equal to
[2021, 18 March Shift-II]

Options:

A. 7

B. 2

C. 5

D. 3

Answer: C

Solution:

Solution:

Given, $f(x) = \frac{x-2}{x-3}$

$g(x) = 2x - 3$

Let $y = f(x) = \frac{x-2}{x-3}$

$\Rightarrow \; xy - 3y = x - 2 \Rightarrow \; xy - x = 3y - 2$

$\Rightarrow \; x(y-1) = 3y - 2$

$\Rightarrow \; x = \frac{3y-2}{y-1}$

$\Rightarrow \; f^{-1}(y) = \frac{3y-2}{y-1}$

$\Rightarrow \; f^{-1}(x) = \frac{3x-2}{x-1}$

Similarly, $g^{-1}(x) = \frac{x+3}{2}$

Given, $f^{-1}(x) + g^{-1}(x) = \frac{13}{2}$

$\Rightarrow \; \frac{3x-2}{x-1} + \frac{x+3}{2} = \frac{13}{2}$

$\Rightarrow \; x^2 + 8x - 7 = 13(x-1)$

$\Rightarrow \; x^2 - 5x + 6 = 0$

$\Rightarrow \; (x-2)(x-3) = 0$

$\Rightarrow \; x = 2, 3$

$\therefore$ Sum $= 2 + 3 = 5$

Question128

The inverse of $y = 5^{\log x}$ is
[2021, 17 March Shift-I]

Options:

A. $x = 5^{\log y}$

B. $x = y^{\log 5}$

C. $x = y^{\frac{1}{\log 5}}$

D. $x = 5^{\frac{1}{\log y}}$

Answer: C

Solution:

Solution:
$y = 5^{\log x}$
Taking log on both sides,
$\Rightarrow \quad \log y = \log x \cdot \log 5$

$\Rightarrow \quad \frac{1}{\log 5} = \frac{\log x}{\log y}$

$\frac{1}{\log 5} = \log_y x$

$x = y^{\frac{1}{\log 5}}$

Question129

Let A = {1, 2, 3, ..., 10} and f : A → A be defined as defined as

$$\mathbf{f(x)} = \begin{cases} x+1 & \text{if } x \text{ is odd} \\ x & \text{if } x \text{ is even} \end{cases}.$$

Then, the number of possible functions g : A → A, such that gof = f

is
[2021, 26 Feb. Shift-II]

Options:

A. 10^5

B. ${}^{10}C_5$

C. 5^5

D. 5 !

Answer: A

Solution:

Solution:

$$f(x) = \begin{cases} x+1 & x \text{is odd} \\ x & x \text{is even.} \end{cases}.$$

Given, $g : A \to A$ such that,
$g(f(x)) = f(x)$
When x is even, then
$g(x) = x$
When x is odd, then
$g(x+1) = x+1$
This implies,
$g(x) = x_1 x$ is even.
$\Rightarrow$ If x is odd, then $g(x)$ can take any value in set A.
So, number of $g(x) = 10^5$

Question130

Let $f, g : N \to N$, such that $f(n+1) = f(n) + f(1)\ \forall n \in N$ and g be any arbitrary function. Which of the following statements is not true?
[2021, 25 Feb. Shift-1]

Options:

A. if $f \circ g$ is one-one, then g is one-one.

B. if f is onto, then $f(n) = n,\ \forall n \in N$.

C. f is one-one.

D. if g is onto, then fog is one-one.

Answer: D

Solution:

Solution:
Given, $f(n+1) = f(n) + f(1),\ \forall n \in N$
$\Rightarrow\ f(n+1) - f(n) = f(1)$
It is an AP with common difference $= f(1)$
Also, general term
$11 = T_n = f(1) + (n-)f(1) = nf(1)$
$\Rightarrow\ f(n) = nf(1)$
Clearly, $f(n)$ is one-one.
For fog to be one-one, g must be one-one.
For f to be onto, $f(n)$ should take all the values of natural numbers.
As, $f(x)$ is increasing, $f(1) = 1$
$\Rightarrow\ f(n) = n$
If g is many-one, then fog is many one.
So, if g is onto, then fog is one-one.

Question131

Let x denote the total number of one-one functions from a set A with 3 elements to a set B with 5 elements and y denote the total number of one-one functions from the set A to the set A × B. Then, [2021, 25 Feb. Shift-II]

Options:

A. $2y = 91x$

B. $2y = 273x$

C. $y = 91x$

D. $y = 273x$

Answer: A

Solution:

Solution:
$x = \{ f : A \to B, f$ is one - one $\}$
$y = \{ g : A \to A \times B, g$ is one one $\}$
Number of elements in $A = 3$ i.e. $|A| = 3$
Similarly, $|B| = 5$
Then, $|A \times B| = |A| \times |B| = 3 \times 5 = 15$
Now, number of one-one function from A to B will be
$1\ {}^5P_3 = \frac{5!}{(5-3)!} = \frac{5!}{2!} = 5 \times 4 \times 3 = 60$
$\therefore\ x = 60$
Now, number of one-one function from A
$1\ \text{to}\ A \times B\ \text{will be}\ = {}^{15}P_3 = \frac{15!}{(15-3)!} = \frac{15!}{12!}$
$= 15 \times 14 \times 13 = 2730$
$\therefore\ y = 2730\ 1c \therefore\ y = 2730$
Thus, $2 \times (2730) = 91 \times (60)$
$\Rightarrow 2y = 91x$

Question132

Let $f : R \to R$ be defined as $f(x) = 2x - 1$ and $g : R - \{1\} \to R$ be defined as $g(x) = \frac{x - \frac{1}{2}}{x - 1}$. Then the composition function $f(g(x))$ is : 24 Feb 2021 Shift 1

Options:

A. onto but not one-one

B. both one-one and onto

C. one-one but not onto

D. neither one-one nor onto

Answer: C

Solution:

Solution:
$f(g(x)) = 2g(x) - 1 = 2\left(\frac{2x-1}{2(x-1)}\right) - 1 = \frac{x}{x-1} = 1 + \frac{1}{x-1}$
Range of $f(g(x)) = mathbb\ R - \{1\}$
Range of $f(g(x))$ is not onto

&f(g(x)) is one-one
So, f(g(x)) is one-one but not onto.

Question133

Let R_1 and R_2 be two relations defined as follows : $R_1 = \{(a, b) \in R^2 : a^2 + b^2 \in Q\}$ and $R_2 = \{(a, b) \in R^2 : a^2 + b^2 \notin Q\}$, where Q is the set of all rational numbers. Then : [Sep. 03, 2020 (II)]

Options:

A. Neither R_1 nor R_2 is transitive.

B. R_2 is transitive but R_1 is not transitive.

C. R_1 is transitive but R_2 is not transitive.

D. R_1 and R_2 are both transitive.

Answer: A

Solution:

Solution:
(a) For R_1 let $a = 1 + \sqrt{2}, b = 1 - \sqrt{2}, c = 8^{1/4}$
$aR_1b \Rightarrow a^2 + b^2 = (1 + \sqrt{2})^2 + (1 - \sqrt{2})^2 = 6 \in Q$
$bR_1c \Rightarrow b^2 + c^2 = (1 - \sqrt{2})^2 + (8^{1/4})^2 = 3 \in Q$
$aR_1c \Rightarrow a^2 + c^2 = (1 + \sqrt{2})^2 + (8^{1/4})^2 = 3 + 4\sqrt{2} \notin Q$
$\therefore R_1$ is not transitive.
For R_2 let $a = 1 + \sqrt{2}, b = \sqrt{2}, c = 1 - \sqrt{2}$
$aR_2b \Rightarrow a^2 + b^2 = (1 + \sqrt{2})^2 + (\sqrt{2})^2 = 5 + 2\sqrt{2} \notin Q$
$bR_2c \Rightarrow b^2 + c^2 = (\sqrt{2})^2 + (1 - \sqrt{2})^2 = 5 - 2\sqrt{2} \notin Q$
$aR_2c \Rightarrow a^2 + c^2 = (1 + \sqrt{2})^2 + (1 - \sqrt{2})^2 = 6 \in Q$
$\therefore R_2$ is not transitive.

Question134

The domain of the function $f(x) = \sin^{-1}\left(\frac{|x|+5}{x^2+1}\right)$ is $(-\infty, -a] \cup [a, \infty]$.

Then a is equal to :
[Sep. 02, 2020 (I)]

Options:

A. $\frac{\sqrt{17}}{2}$

B. $\frac{\sqrt{17}-1}{2}$

C. $\frac{1+\sqrt{17}}{2}$

D. $\frac{\sqrt{17}}{2}+1$

Answer: C

Solution:

Solution:

$\because f(x) = \sin^{-1}\left(\frac{|x|+5}{x^2+1}\right)$

$\therefore -1 \le \frac{|x|+5}{x^2+1} \le 1$

$\Rightarrow |x| + 5 \le x^2 + 1$

$[\because x^2 + 1 \ne 0]$

$\Rightarrow x^2 - |x| - 4 \ge 0$

$\Rightarrow \left(|x| - \frac{1-\sqrt{17}}{2}\right)\left(|x| - \frac{1+\sqrt{17}}{2}\right) \ge 0$

$\Rightarrow x \in \left(-\infty, -\frac{1+\sqrt{17}}{2}\right) \cup \left[\frac{1+\sqrt{17}}{2}, \infty\right)$

$\therefore a = \frac{1+\sqrt{17}}{2}$

Question135

If $R = \{(x, y) : x, y \in Z, x^2 + 3y^2 \le 8\}$ is a relation on the set of integers Z, then the domain of R^{-1} is :
[Sep. 02, 2020 (I)]

Options:

A. $\{-2, -1, 1, 2\}$

B. $\{0, 1\}$

C. $\{-2, -1, 0, 1, 2\}$

D. $\{-1, 0, 1\}$

Answer: D

Solution:

Solution:
Since, $R = \{(x, y) : x, y \in Z, x^2 + 3y^2 \leq 8\}$
$\therefore R = \{(1, 1), (2, 1), (1, -1), (0, 1), (1, 0)\}$
$\Rightarrow D_{R^{-1}} = \{-1, 0, 1\}$

Question136

Let [t] denote the greatest integer ≤t. Then the equation in x, $[x]^2 + 2[x + 2] - 7 = 0$ has : [Sep. 04, 2020 (I)]

Options:

A. exactly two solutions

B. exactly four integral solutions

C. no integral solution

D. infinitely many solutions

Answer: D

Solution:

Solution:
The given equation $[x]^2 + 2[x] + 4 - 7 = 0$
$\Rightarrow [x]^2 + 2[x] - 3 = 0$
$\Rightarrow [x]^2 + 3[x] - [x] - 3 = 0$

$\Rightarrow([x]+3)([x]-1)=0 \Rightarrow [x]=1$ or -3 $\Rightarrow x \in [-3,-2) \cup [1,2)$
$\therefore$ The equation has infinitely many solutions.

Question137

Let f (x) be a quadratic polynomial such that $f(-1)+f(2)=0$. If one of the roots of f (x) = 0 is 3, then its other root lies in: [Sep. 02, 2020 (II)]

Options:

A. (-1,0)

B. (1,3)

C. (-3,-1)

D. (0,1)

Answer: A

Solution:

Solution:
Let $f(x)=ax^2+bx+c$
Given : $f(-1)+f(2)=0$
$a-b+c+4a+2b+c=0$
$\Rightarrow 5a+b+2c=0$ .....(i)
and $f(3)=0 \Rightarrow 9a+3b+c=0$ .....(ii)
From equations (i) and (ii),
$\frac{a}{1-6}=\frac{b}{18-5}=\frac{c}{15-9} \Rightarrow \frac{a}{-5}=\frac{b}{13}=\frac{c}{6}$
Product of roots, $\alpha\beta=\frac{c}{a}=\frac{-6}{5}$ and $\alpha=3$
$\Rightarrow \beta=\frac{-2}{5} \in (-1,0)$

Question138

Let $f(1,3) \rightarrow R$ be a function defined by $f(x)=\frac{x[x]}{1+x^2}$ where [x] denotes the greatest integer $\leq x$. Then the range of f is: [Jan. 8, 2020 (II)]

Options:

A. $\left(\frac{2}{5}, \frac{3}{5}\right) \cup \left(\frac{3}{4}, \frac{4}{5}\right)$

B. $\left(\frac{2}{5}, \frac{1}{2}\right) \cup \left(\frac{3}{5}, \frac{4}{5}\right)$

C. $\left(\frac{2}{5}, \frac{4}{5}\right)$

D. $\left(\frac{3}{5}, \frac{4}{5}\right)$

Answer: B

Solution:

Solution:

$$f(x) \begin{cases} \frac{x}{x^2+1} & x \in (1, 2) \\ \frac{2x}{x^2+1} & x \in [2, 3). \end{cases}$$

$$f'(x) \begin{cases} \frac{1-x^2}{1+x^2} & x \in (1, 2) \\ \frac{1-2x^2}{1+x^2} & x \in [2, 3). \end{cases}$$

$\therefore f(x)$ is a decreasing function

$\therefore \ y \in \left(\frac{2}{5}, \frac{1}{2}\right) \cup \left(\frac{6}{10}, \frac{4}{5}\right]$

$\Rightarrow \ y \in \left(\frac{2}{5}, \frac{1}{2}\right) \cup \left(\frac{3}{5}, \frac{4}{5}\right]$

Question139

Let A = {a, b, c} and B = {1, 2, 3, 4}. Then the number of elements in the set C = { f : A → B | 2 ∈ f (A) and f is not one-one } is _____.

[NA Sep. 05,2020 (II)]

Answer: 19

Solution:

Solution:
The desired functions will contain either one element or two elements in its codomain of which '2' always belongs to $f(A)$.
$\therefore$ The set B can be $\{2\}, \{1, 2\}, \{2, 3\}, \{2, 4\}$
Total number of functions $= 1 + (2^3 - 2)3 = 19$

Question140

The inverse function of $f(x) = \frac{8^{2x} - 8^{-2x}}{8^{2x} + 8^{-2x}}$, $x \in (-1, 1)$, is _____.

[Jan. 8, 2020 (I)]

Options:

A. $\frac{1}{4}\log_e\left(\frac{1+x}{1-x}\right)$

B. $\frac{1}{4}(\log_8 e)\log_e\left(\frac{1-x}{1+x}\right)$

C. $\frac{1}{4}\log_e\left(\frac{1-x}{1+x}\right)$

D. $\frac{1}{4}(\log_8 e)\log_e\left(\frac{1+x}{1-x}\right)$

Answer: A

Solution:

Solution:

$$y = \frac{8^{2x} - 8^{-2x}}{8^{2x} + 8^{-2x}}$$

$$\frac{1+y}{1-y} = \frac{8^{2x}}{8^{-2x}} \Rightarrow 8^{4x} = \frac{1+y}{1-y}$$

$$\Rightarrow 4x = \log_8\left(\frac{1+y}{1-y}\right)$$

$$\Rightarrow x = \frac{1}{4}\log_8\left(\frac{1+y}{1-y}\right)$$

$$\therefore f^{-1}(x) = \frac{1}{4}\log_8\left(\frac{1+x}{1-x}\right)$$

Question141

If $g(x) = x^2 + x - 1$ and $(gof)(x) = 4x^2 - 10x + 5$, then $f\left(\frac{5}{4}\right)$ is equal to: [Jan. 7, 2020 (I)]

Options:

A. $\frac{3}{2}$

B. $-\frac{1}{2}$

C. $\frac{1}{2}$

D. $-\frac{3}{2}$

Answer: B

Solution:

Solution:

$$(gof)(x) = g(f(x)) = f^2(x) + f(x) - 1$$

$$g\left(f\left(\frac{5}{4}\right)\right) = 4\left(\frac{5}{4}\right)^2 - 10 \cdot \frac{5}{4} + 5 = -\frac{5}{4}$$

$$[\because g(f(x)) = 4x^2 - 10x + 5]$$

$$g\left(f\left(\frac{5}{4}\right)\right) = f^2\left(\frac{5}{4}\right) + f\left(\frac{5}{4}\right) - 1$$

$$-\frac{5}{4} = f^2\left(\frac{5}{4}\right) + f\left(\frac{5}{4}\right) - 1$$

$$f^2\left(\frac{5}{4}\right) + f\left(\frac{5}{4}\right) + \frac{1}{4} = 0$$

$\left(f\left(\frac{5}{4}\right)+\frac{1}{2}\right)^2=0$

$t\left(\frac{5}{4}\right)=-\frac{1}{2}$

Question142

For a suitably chosen real constant a, let a function,$f: R-\{-a\} \to R$ be defined by $f(x)=\frac{a-x}{a+x}$. Further suppose that for any real number $x \neq -a$ and $f(x) \neq -a$, $(fof)(x)=x$. Then $f\left(-\frac{1}{2}\right)$ is equal to:

[Sep. 06, 2020 (II)]

Options:

A. $\frac{1}{3}$

B. $-\frac{1}{3}$

C. -3

D. 3

Answer: D

Solution:

Solution:

$$f(f(x))=\frac{a-\left(\frac{a-x}{a+x}\right)}{a+\left(\frac{a-x}{a+x}\right)}=x$$

$$\Rightarrow \frac{a-ax}{1+x}=f(x) \Rightarrow \frac{a(1-x)}{1+x}=\frac{a-x}{a+x} \Rightarrow a=1$$

$$\therefore f(x)=\frac{1-x}{1+x} \Rightarrow f\left(-\frac{1}{2}\right)=3$$

Question143

Let $f : R \to R$ be defined by $f(x) = \frac{x}{1+x^2}$, $x \in R$. Then the range of f is :
[Jan. 11, 2019 (I)]

Options:

A. $\left[-\frac{1}{2}, \frac{1}{2}\right]$

B. $R - [-1, 1]$

C. $R - \left[-\frac{1}{2}, \frac{1}{2}\right]$

D. $(-1, 1) - \{0\}$

Answer: A

Solution:

Solution:

$f(x) = \frac{x}{1+x^2}, x \in R$

Let, $y = \frac{x}{1+x^2}$

$\Rightarrow yx^2 - x + y = 0 \Rightarrow x = \frac{1 \pm \sqrt{1-4y^2}}{2}$

$\Rightarrow 1 - 4y^2 \geq 0$

$\Rightarrow 1 \geq 4y^2$

$\Rightarrow |y| \leq \frac{1}{2}$

$\Rightarrow -\frac{1}{2} \leq y \leq \frac{1}{2}$

$\Rightarrow$ The range of f is $\left[-\frac{1}{2}, \frac{1}{2}\right]$

Question144

The domain of the definition of the function
$f(x) = \frac{1}{4-x^2} + \log_{10}(x^3 - x)$ is:
[April. 09, 2019 (II)]

Options:

A. $(-1,0)\cup(1,2)\cup(3,\infty)$

B. $(-2,-1)\cup(-1,0)\cup(2,\infty)$

C. $(-1,0)\cup(1,2)\cup(2,\infty)$

D. $(1,2)\cup(2,\infty)$

Answer: C

Solution:

Solution:
To determine domain, denominator $\neq 0$ and $x^3 - x > 0$
i.e., $4 - x^2 \neq 0 x \neq \pm 2$ .....(1)
and $x(x-1)(x+1) > 0$



$x \in (-1, 0) \cup (1, \infty)$ .....(2)
Hence domain is intersection of (1)&(2).
i.e., $x \in (-1, 0) \cup (1, 2) \cup (2, \infty)$

Question145

If $f(x) = \log_e\left(\frac{1-x}{1+x}\right), |x|<1$, then $f\left(\frac{2x}{1+x^2}\right)$ is equal to

[April 8, 2019 (I)]

Options:

A. $2f(x)$

B. $2f(x^2)$

C. $(f(x))^2$

D. $-2f(x)$

Answer: A

Solution:

Solution:

$f(x)=\log\left(\frac{1-x}{1+x}\right),|x|<1$

$$f\left(\frac{2x}{1+x^2}\right)=\log\left(\frac{1-\frac{2x}{1+x^2}}{1+\frac{2x}{1+2x^2}}\right)$$

$$=\log\left(\frac{1+x^2-2x}{1+x^2+2x}\right)=\log\left(\frac{1-x}{1+x}\right)^2$$

$$=2\log\left(\frac{1-x}{1+x}\right)=2f(x)$$

Question146

Let $f(x)=a^x(a>0)$ be written as $f(x)=f_1(x)+f_2(x)$, where $f_1(x)$ is an even function and $f_2(x)$ is an odd function. Then $f_1(x+y)+f_1(x-y)$ equals:
[April. 08, 2019 (II)]

Options:

A. $2f_1(x)f_1(y)$

B. $2f_1(x+y)f_1(x-y)$

C. $2f_1(x)f_2(y)$

D. $2f_1(x+y)f_2(x-y)$

Answer: A

Solution:

Solution:
Given function can be written as

$$f(x)=a^x=\left(\frac{a^x+a^{-x}}{2}\right)+\left(\frac{a^x-a^{-x}}{2}\right)$$

where $f_1(x)=\frac{a^x+a^{-x}}{2}$ is even function

$f_2(x)=\frac{a^x-a^{-x}}{2}$ is odd function

$\Rightarrow f_1(x+y)+f_1(x-y)$

$$= \left(\frac{a^{x+y}+a^{-x-y}}{2}\right)+\left(\frac{a^{x-y}+a^{-x+y}}{2}\right)$$

$$= \frac{1}{2}[a^x(a^y+a^{-y})+a^{-x}(a^y+a^{-y})]$$

$$= \frac{(a^x+a^{-x})(a^y+a^{-y})}{2} = 2f_1(x)\cdot f_1(y)$$

Question147

Let a function $f : (0, \infty) \to (0, \infty)$ be defined by $f(x) = \left|1-\frac{1}{x}\right|$. Then f is :
[Jan. 11, 2019 (II)]

Options:

A. not injective but it is surjective

B. injective only

C. neither injective nor surjective

D. (Bonus)

Answer: D

Solution:

Solution:
$f : (0, \infty) \to (0, \infty)$
$f(x) = \left|1-\frac{1}{x}\right|$ is not a function
$\because f(1) = 0$ and $1 \in$ domain but $0 \notin$ co-domain
Hence, $f(x)$ is not a function.

Question148

The number of functions f from {1, 2, 3, ..., 20} onto {1, 2, 3, ..., 20} such that f (k) is a multiple of 3, whenever k is a multiple of 4 is :
[Jan. 11, 2019 (II)]

Options:

A. $6^5 \times (15)!$

B. $5! \times 6!$

C. $(15)! \times 6!$

D. $5^6 \times 15$

Answer: C

Solution:

Solution:
Domain and codomain $= \{1, 2, 3, ..., 20\}$.
There are five multiple of 4 as 4,8,12,16 and 20 .
and there are 6 multiple of 3 as 3,6,9,12,15,18 .
Since, when ever k is multiple of 4 then $f(k)$ is multiple of 3 then total number of arrangement $= {}^6c_5 \times 5! = 6!$
Remaining 15 elements can be arranged in $15!$ ways.
Since, for every input, there is an output
$\Rightarrow$ function $f(k)$ in onto
$\therefore$ Total number of arrangement $= 15!.6!$

Question149

Let N be the set of natural numbers and two functionsf and g be

defined as f, g : N → N such that f (n) = $\begin{cases} \frac{n+1}{2} & \text{if n is odd} \\ \frac{n}{2} & \text{if n is even} \end{cases}$

and g(n) = n − (−1)^n. Then f og is:
[Jan. 10, 2019 (II)]

Options:

A. onto but not one-one.

B. one-one but not onto.

C. both one-one and onto.

D. neither one-one nor onto.

Answer: A

Solution:

Solution:

$$f(n) = \begin{cases} \frac{n+1}{2}, & \text{if } n \text{ is odd} \\ \frac{n}{2}, & \text{if } n \text{ is even} \end{cases}$$

$$g(n) = \begin{cases} 2, & n = 1 \\ 1, & n = 2 \\ 4, & n = 3 \\ 3, & n = 4 \\ 6, & n = 5 \\ 5, & n = 6 \end{cases}$$

Then,

$$f(g(n)) = \begin{cases} \frac{g(n)+1}{2}, & \text{if } g(n) \text{ is odd} \\ \frac{g(n)}{2}, & \text{if } g(n) \text{ is even} \end{cases}$$

$$f(g(n)) = \begin{cases} 1, & n = 1 \\ 1, & n = 2 \\ 2, & n = 3 \\ 2, & n = 4 \\ 3, & n = 5 \\ 3, & n = 6 \\ : & : \\ : & : \\ : & : \end{cases}$$

⇒ fog is onto but not one - one

Question150

Let A = { x ∈ R : x is not a positive integer }. Define a function $f : A \to R$ as $f(x) = \frac{2x}{x-1}$, then f is:
[Jan. 09, 2019 (II)]

Options:

A. not injective

B. neither injective nor surjective

C. surjective but not injective

D. injective but not surjective

Answer: D

Solution:

Solution:
As $A = \{x \in R : x \text{ is not a positive integer}\}$
A function $f : A \to R$ given by $f(x) = \frac{2x}{x-1}$
$f(x_1) = f(x_2) \Leftrightarrow x_1 = x_2$
So, f is one-one.
As $f(x) \neq 2$ for any $x \in A \Rightarrow f$ is not onto.
Hence f is injective but not surjective.

Question151

For $x \in \left(0, \frac{3}{2}\right)$, let $f(x) = \sqrt{x}$, $g(x) = \tan x$ and $h(x) = \frac{1-x^2}{1+x^2}$

If $\phi(x) = ((hof)og)(x)$, then $\phi\left(\frac{\pi}{3}\right)$ is equal to

[April 12, 2019 (I)]

Options:

A. $\tan\frac{\pi}{12}$

B. $\tan\frac{11\pi}{12}$

C. $\tan\frac{7\pi}{12}$

D. $\tan\frac{5\pi}{12}$

Answer: B

Solution:

Solution:

$\because \phi(x) = (\,(\text{ hof }) \text{ og }\,)(x)$

$\because \phi\left(\frac{\pi}{3}\right) = h\left(f\left(g\left(\frac{\pi}{3}\right)\right)\right) = h(f(\sqrt{3})) = h(3^{1/4})$

$= \frac{1-\sqrt{3}}{1+\sqrt{3}} = -\frac{1}{2}(1+3-2\sqrt{3}) = \sqrt{3}-2 = -(-\sqrt{3}+2)$

$= -\tan 15° = \tan(180° - 15°) = \tan\left(\pi - \frac{\pi}{12}\right) = \tan\frac{11\pi}{12}$

Question152

Let $f(x) = x^2$, $x \in R$. For any $A \subseteq R$, define $g(A) = \{x \in R : f(x) \in A\}$. If $S = [0, 4]$, then which one of the following statements is not true?
[April 10, 2019 (I)]

Options:

A. $g(f(S)) \neq S$

B. $f(g(S)) = S$

C. $g(f(S)) = g(S)$

D. $f(g(S)) \neq f(S)$

Answer: C

Solution:

Solution:

$f(x) = x^2;\ x \in R$

$g(A) = \{x \in R : f(x) \in A\} S = [0, 4]$

$g(S) = \{x \in R : f(x) \in S\}$

$= \{x \in R : 0 \leq x^2 \leq 4\} = \{x \in R : -2 \leq x \leq 2\}$

$\therefore g(S) \neq S \therefore f(g(S)) \neq f(S)$

$g(f(S)) = \{x \in R : f(x) \in f(S)\}$

$= \{x \in R : x^2 \in S^2\} = \{x \in R : 0 \leq x^2 \leq 16\}$

$= \{x \in R : -4 \leq x \leq 4\}$

$\therefore g(f(S)) \neq g(S)$

$\therefore g(f(S)) = g(S)$ is incorrect.

Question153

For $x \in R - \{0, 1\}$, let $f_1(x) = \frac{1}{x}$, $f_2(x) = 1 - x$ and $f_3(x) = \frac{1}{1-x}$ be three given functions. If a function, J (x) satisfies $(f_2 o J o f_1)(x) = f_3(x)$ then J (x) is equal to:
[Jan. 09, 2019 (I)]

Options:

A. $f_3(x)$

B. $\frac{1}{x} f_3(x)$

C. $f_2(x)$

D. $f_1(x)$

Answer: A

Solution:

Solution:
The given relation is

$$(f_2 o J \text{ of}_1)(x) = f_3(x) = \frac{1}{1-x}$$

$$\Rightarrow (f_2 J)(f_1(x)) = \frac{1}{1-x}$$

$$\Rightarrow (f_2 o J)\left(\frac{1}{x}\right) = \frac{1}{1-\frac{1}{\frac{1}{x}}} = \frac{\frac{1}{x}}{\frac{1}{x}-1} \left[\because f_1(x) = \frac{1}{x}\right]$$

$$\Rightarrow (f_2 o J)(x) = \frac{x}{x-1} \left[\frac{1}{x} \text{ is replaced by } x\right]$$

$$\Rightarrow f_2(J(x)) = \frac{x}{x-1}$$

$$\Rightarrow 1 - J(x) = \frac{x}{x-1} [\because f_2(x) = 1 - x]$$

$$\therefore J(x) = 1 - \frac{x}{x-1} = \frac{1}{1-x} = f_3(x)$$

Question154

Let N denote the set of all natural numbers. Define two binary relations on N as $R_1 = \{(x, y) \in N \times N : 2x + y = 10\}$ and

$R_2 = \{(x, y) \in N \times N : x + 2y = 10\}$. Then
[Online April 16, 2018]

Options:

A. Both R_1 and R_2 are transitive relations

B. Both R_1 and R_2 are symmetric relations

C. Range of R_2 is $\{1, 2, 3, 4\}$

D. Range of R_1 is $\{2, 4, 8\}$

Answer: C

Solution:

Solution:
Here, $R_1 = \{(x, y) \in N \times N : 2x + y = 10\}$ and
$R_2 = \{(x, y) \in N \times N : x + 2y = 10\}$
For R_1; $2x + y = 10$ and $x, y \in N$
So, possible values for x and y are:
$x = 1, y = 8$ i.e. (1,8);
$x = 2, y = 6$ i.e. (2,6);
$x = 3, y = 4$ i.e. (3,4) and $x = 4, y = 2$ i.e. (4,2).
$R_1 = \{(1, 8), (2, 6), (3, 4), (4, 2)\}$
Therefore, Range of R_1 is $\{2, 4, 6, 8\}$
R_1 is not symmetric
Also, R_1 is not transitive because $(3, 4), (4, 2) \in R_1$ but $(3,2) \notin R_1$
Thus, options A, B and D are incorrect.
For R_2; $x + 2y = 10$ and $x, y \in N$
So, possible values for x and y are:$x = 8, y = 1$ i.e. (8,1);
$x = 6, y = 2$ i.e. (6,2);
$x = 4, y = 3$ i.e. (4,3) and
$x = 2, y = 4$ i.e. (2, 4)
$R_2 = \{(8, 1), (6, 2), (4, 3), (2, 4)\}$
Therefore, Range of R_2 is $\{1, 2, 3, 4\}$
R_2 is not symmetric and transitive.

Question155

Consider the following two binary relations on the set $A = \{a, b, c\}$: $R_1 = \{(c, a)(b, b), (a, c), (c, c), (b, c), (a, a)\}$and $R_2 = \{(a, b), (b, a), (c, c), (c, a), (a, a), (b, b), (a, c)$. Then
[Online April 15, 2018]

Options:

A. R_2 is symmetric but it is not transitive

B. Both R_1 and R_2 are transitive

C. Both R_1 and R_2 are not symmetric

D. R_1 is not symmetric but it is transitive

Answer: A

Solution:

Solution:
Both R_1 and R_2 are symmetric as
For any $(x, y) \in R_1$, we have
$(y, x) \in R_1$ and similarly for R_2
Now, for R_2, $(b, a) \in R_2$, $(a, c) \in R_2$ but $(b, c) \notin R_2$
Similarly, for R_1, $(b, c) \in R_1$, $(c, a) \in R_1$ but $(b, a) \notin R_1$
Therefore, neither R_1 nor R_2 is transitive.

Question156

Let $f : A \to B$ be a function defined as $f(x) = \frac{x-1}{x-2}$, where $A = R - \{2\}$ and $B = R - \{1\}$. Then f is
[Online April 15, 2018]

Options:

A. invertible and $f^{-1}(y) = \frac{2y+1}{y-1}$

B. invertible and $f^{-1}(y) = \frac{3y-1}{y-1}$

C. no invertible

D. invertible and $f^{-1}(y) = \frac{2y-1}{y-1}$

Answer: D

Solution:

Solution:
Suppose $y = f(x)$
$\Rightarrow y = \frac{x-1}{x-2}$
$\Rightarrow yx - 2y = x - 1$
$\Rightarrow (y-1)x = 2y - 1$
$\Rightarrow x = f^{-1}(y) = \frac{2y-1}{y-1}$
As the function is invertible on the given domain and its inverse can be obtained as above.

Question157

The function $f : R \rightarrow \left[-\frac{1}{2}, \frac{1}{2}\right]$ defined as $f(x) = \frac{x}{1+x^2}$, is: [2017]

Options:

A. neither injective nor surjective

B. invertible

C. injective but not surjective

D. surjective but not injective

Answer: D

Solution:

Solution:
We have $f : R \rightarrow \left[-\frac{1}{2}, \frac{1}{2}\right]$,
$f(x) = \frac{x}{1+x^2} \; \forall x \in R$
$\Rightarrow f'(x) = \frac{(1+x^2).1 - x.2x}{(1+x^2)^2} = \frac{-(x+1)(x-1)}{(1+x^2)^2}$

sign of $f'(x)$
$\Rightarrow f'(x)$ changes sign in different intervals.
$\therefore$ Not injective Now $y = \frac{x}{1+x^2}$
$\Rightarrow y + yx^2 = x$
$\Rightarrow yx^2 - x + y = 0$
For $y \neq 0, D = 1 - 4y^2 \geq 0$
$\Rightarrow y \in \left[\frac{-1}{2}, \frac{1}{2}\right] - \{0\}$
For $y = 0 \Rightarrow x = 0$
$\therefore$ Range is $\left[\frac{-1}{2}, \frac{1}{2}\right]$
$\Rightarrow$ Surjective but not injective

Question158

The function $f : N \rightarrow N$ defined by $f(x) = x - 5\left[\frac{x}{5}\right]$, where N is set of natural numbers and [x] denotes the greatest integer less than or equal to x, is:
[Online April 9, 2017]

Options:

A. one-one and onto.

B. one-one but not onto.

C. onto but not one-one.

D. neither one-one nor onto.

Answer: D

Solution:

Solution:
$\left.\begin{array}{l} f(1) = 1 - 5[1/5] = 1 \\ f(6) = 6 - 5[6/5] = 1 \end{array}\right\} \rightarrow$ Many one
$f(10) = 10–5(2) = 0$ which is not in co–domain.
Neither one–one nor onto.

Question159

Let $f(x) = 2^{10} . x + 1$ and $g(x) = 3^{10} . x - 1$. If $(f \circ g)(x) = x$, then x is equal to:
[Online April 8, 2017]

Options:

A. $\frac{3^{10} - 1}{3^{10} - 2^{-10}}$

B. $\frac{2^{10} - 1}{2^{10} - 3^{-10}}$

C. $\frac{1 - 3^{-10}}{2^{10} - 3^{-10}}$

D. $\frac{1 - 2^{-10}}{3^{10} - 2^{-10}}$

Answer: D

Solution:

Solution:
$f(g(x)) = x$
$\Rightarrow f(3^{10}x - 1) = 2^{10}(3^{10} . x - 1) + 1 = x$
$\Rightarrow 2^{10}(3^{10}x - 1) + 1 = x$
$\Rightarrow x(6^{10} - 1) = 2^{10} - 1$
$\Rightarrow x = \frac{2^{10} - 1}{6^{10} - 1} = \frac{1 - 2^{-10}}{3^{10} - 2^{-10}}$

Question160

For $x \in R, x \neq 0$, let $f_0(x) = \frac{1}{1 - x}$ and $f_{n+1}(x) = f_0(f_n(x))n = 0, 1, 2, ...$

Then the value of $f_{100}(3) + f_1\left(\frac{2}{3}\right) + f_2\left(\frac{3}{2}\right)$ is equal to :

[Online April 9, 2016]

Options:

A. $\frac{8}{3}$

B. $\frac{4}{3}$

C. $\frac{5}{3}$

D. $\frac{1}{3}$

Answer: C

Solution:

Solution:

$f_1(x) = f_{0+1}(x) = f_0(f_0(x)) = \frac{1}{1-\frac{1}{1-x}} = \frac{x-1}{x}$

$f_2(x) = f_{1+1}(x) = f_0(f_1(x)) = \frac{1}{1-\frac{x-1}{x}} = x$

$f_3(x) = f_{2+1}(x) = f_0(f_2(x)) = f_0(x) = \frac{1}{1-x}$

$f_4(x) = f_{3+1}(x) = f_0(f_3(x)) = \frac{x-1}{x}$

$\therefore f_0 = f_3 = f_6 = ... = \frac{1}{1-x}$

$f_1 = f_4 = f_7 = ... = \frac{x-1}{x}$

$f_2 = f_5 = f_8 = ... = x$

$f_{100}(3) = \frac{3-1}{3} = \frac{2}{3} f_1\left(\frac{2}{3}\right) = \frac{\frac{2}{3}-1}{\frac{2}{3}} = -\frac{1}{2}$

$f_2\left(\frac{3}{2}\right) = \frac{3}{2}$

$\therefore f_{100}(3) + f_1\left(\frac{2}{3}\right) + f_2\left(\frac{3}{2}\right) = \frac{5}{3}$

Question161

Let A = {x_1, x_2, ..., x_7} and B = {y_1, y_2, y_3} be two sets containing seven and three distinct elements respectively. Then the total number of functions f : A → B that are onto, if there exist exactly three elements x in A such that f (x) = y_2, is equal to

(Online April 11, 2015)

Options:

A. $14.{}^{7}C_{3}$

B. $16.{}^{7}C_{3}$

C. $14.{}^{7}C_{2}$

D. $12..{}^{7}C_{2}$

Answer: A

Solution:

Solution:
Number of onto function such that exactly three elements in $x \in A$ such that $f(x) = \frac{1}{2}$ is equal to
$= {}^{7}C_{3}, \{2^{4} - 2\} = 14 . {}^{7}C_{3}$

Question162

Let $f : R \rightarrow R$ be defined by $f(x) = \frac{|x| - 1}{|x| + 1}$ then f is: [Online April 19, 2014]

Options:

A. both one-one and onto

B. one-one but not onto

C. onto but not one-one

D. neither one-one nor onto.

Answer: C

Solution:

Solution:
$f(x) = \frac{|x| - 1}{|x| + 1}$
for one-one function if $f(x_1) = f(x_2)$ then
x_1 must be equal to x_2
Let $f(x_1) = f(x_2)$

$\frac{|x_1|-1}{|x_1|+1}=\frac{|x_2|-1}{|x_2|+1}$ $|x_1||x_2|+|x_1|-|x_2|-1=|x_1||x_2|-|x_1|+|x_2|-1$

$\Rightarrow |x_1|-|x_2|=|x_2|-|x_1|$

$2|x_1|=2|x_2|$

$|x_1|=|x_2|$

$x_1=x_2, x_1=-x_2$

here x_1 has two values therefore function is many one function.

For onto : $f(x)=\frac{|x|-1}{|x|+1}$

for every value of $f(x)$ there is a value of x in domain set.

If $f(x)$ is negative then $x=0$

for all positive value of $f(x)$, domain contain atleast one element. Hence $f(x)$ is onto function.

Question163

Let P be the relation defined on the set of all real numbers such that $P = \{(a, b) : \sec^2 a - \tan^2 b = 1\}$. Then P is: [Online April 9, 2014]

Options:

A. reflexive and symmetric but not transitive.

B. reflexive and transitive but not symmetric.

C. symmetric and transitive but not reflexive.

D. an equivalence relation.

Answer: D

Solution:

Solution:

$P = \{(a, b) : \sec^2 a - \tan^2 b = 1\}$

For reflexive :

$\sec^2 a - \tan^2 a = 1$(true $\forall a$)

For symmetric :

$\sec^2 b - \tan^2 a = 1$

L.H.S

$1+\tan^2 b-(\sec^2 a-1)=1+\tan^2 b-\sec^2 a+1$

$=-(\sec^2 a-\tan^2 b)+2$

$=-1+2=1$

So, Relation is symmetric For transitive :

if $\sec^2 a - \tan^2 b = 1$ and $\sec^2 b - \tan^2 c = 1$

$\sec^2 a - \tan^2 c = (1 + \tan^2 b) - (\sec^2 b - 1)$
$= -\sec^2 b + \tan^2 b + 2$
$= -1 + 2 = 1$
So, Relation is transitive.
Hence, Relation P is an equivalence relation

Question164

Let $f(n) = \left[\frac{1}{3} + \frac{3n}{100} \right] n$, where [n] denotes the greatest integer less than or equal to n. Then $\sum_{n=1}^{56} f(n)$ is equal to:
[Online April 19, 2014]

Options:

A. 56

B. 689

C. 1287

D. 1399

Answer: D

Solution:

Solution:
Let $f(n) = \left[\frac{1}{3} + \frac{3n}{100} \right] n$
where $[n]$ is greatest integer function,
$= \left[0.33 + \frac{3n}{100} \right] n$
For $n = 1, 2, \ldots, 22$, we get $f(n) = 0$ and for $n = 23, 24, \ldots, 55$, we get $f(n) = 1 \times n$ For $n = 56$, $f(n) = 2 \times n$
So, $\sum_{n=1}^{56} f(n) = 1(23) + 1(24) + \ldots + 1(55) + 2(56)$
$= (23 + 24 + \ldots + 55) + 112$
$= \frac{33}{2}[46 + 32] + 112$
$= \frac{33}{2}(78) + 112 = 1399$

Question165

Let f be an odd function defined on the set of real numbers such that for $x \geq 0$, $f(x) = 3\sin x + 4\cos x$ Then f (x) at $x = -\frac{11\pi}{6}$ is equal to: [Online April 11, 2014]

Options:

A. $\frac{3}{2} + 2\sqrt{3}$

B. $-\frac{3}{2} + 2\sqrt{3}$

C. $\frac{3}{2} - 2\sqrt{3}$

D. $-\frac{3}{2} - 2\sqrt{3}$

Answer: C

Solution:

Solution:
Given f be an odd function
$f(x) = 3\sin x + 4\cos x$

Now, $f\left(\frac{-11\pi}{6}\right) = 3\sin\left(\frac{-11\pi}{6}\right) + 4\cos\left(\frac{-11\pi}{6}\right)$

$f\left(\frac{-11\pi}{6}\right) = 3\sin\left(-2\pi + \frac{\pi}{6}\right) + 4\cos\left(-2\pi + \frac{\pi}{6}\right)$

$f\left(\frac{-11\pi}{6}\right) = 3\sin\left\{-\left(2\pi - \frac{\pi}{6}\right)\right\} + 4\cos\left\{-\left(2\pi - \frac{\pi}{6}\right)\right\}$

$$\left\{\begin{array}{c}\text{For odd functions} \\ \sin(-\theta) = -\sin\theta \\ \text{and } \cos(-\theta) = \cos\theta\end{array}\right\}$$

$\therefore f\left(\frac{-11\pi}{6}\right) = -3\sin\left(2\pi - \frac{\pi}{6}\right) - 4\cos\left(2\pi - \frac{\pi}{6}\right)$

$\Rightarrow f\left(\frac{-11\pi}{6}\right) = +3\sin\left(\frac{\pi}{6}\right) - 4\cos\frac{\pi}{6}$

$\Rightarrow f\left(\frac{-11\pi}{6}\right) = 3\times\frac{1}{2} - 4\times\frac{\sqrt{3}}{2}$

or $f\left(\frac{-11\pi}{6}\right) = \frac{3}{2} - 2\sqrt{3}$

Question166

If g is the inverse of a function f and $f'(x) = \frac{1}{1+x^5}$, then $g'(x)$ is equal to:
[2014]

Options:

A. $\frac{1}{1+\{g(x)\}^5}$

B. $1+\{g(x)\}^5$

C. $1+x^5$

D. $5x^4$

Answer: B

Solution:

Solution:
Since $f(x)$ and $g(x)$ are inverse of each other
$\therefore g'(f(x)) = \frac{1}{f'(x)}$
$\Rightarrow g'(f(x)) = 1+x^5 \quad \left(\because f'(x) = \frac{1}{1+x^5}\right)$
Here $x = g(y)$
$\therefore g'(y) = 1+[g(y)]^5$
$\Rightarrow g'(x) = 1+(g(x))^5$

Question167

Let $R = \{(x, y) : x, y \in N$. and $x^2 - 4xy + 3y^2 = 0\}$, where N is the set of all natural numbers. Then the relation R is:
[Online April 23, 2013]

Options:

A. reflexive but neither symmetric nor transitive.

B. symmetric and transitive.

C. reflexive and symmetric,

D. reflexive and transitive.

Answer: D

Solution:

Solution:
$R = \{(x, y) : x, y \in N$ and $x^2 - 4xy + 3y^2 = 0\}$
Now, $x^2 - 4xy + 3y^2 = 0$
$\Rightarrow (x - y)(x - 3y) = 0$
$\therefore x = y$ or $x = 3y$
$\therefore R = \{(1, 1), (3, 1), (2, 2), (6, 2), (3, 3)(9, 3), \ldots\ldots\}$
Since $(1, 1), (2, 2), (3, 3), \ldots\ldots$ are present in the relation, therefore R is reflexive.
Since (3,1) is an element of R but (1,3) is not the element of R, therefore R is not symmetric
Here $(3,1) \in R$ and $(1,1) \in R \Rightarrow (3, 1) \in R$ $(6,2) \in R$ and $(2,2) \in R \Rightarrow (6, 2) \in R$
For all such $(a, b) \in R$ and $(b, c) \in R$
$\Rightarrow (a, c) \in R$
Hence R is transitive.

Question168

Let R = {(3, 3)(5, 5), (9, 9), (12, 12), (5, 12), (3, 9), (3, 12), (3, 5)} be a relation on the set A = {3, 5, 9, 12}. Then, R is : [Online April 22, 2013]

Options:

A. reflexive, symmetric but not transitive.

B. symmetric, transitive but not reflexive.

C. an equivalence relation.

D. reflexive, transitive but not symmetric.

Answer: D

Solution:

Solution:
Let $R = \{(3, 3), (5, 5), (9, 9), (12, 12), (5, 12), (3, 9), (3, 12),$ (3,5)\} be a relation on set
$A = \{3, 5, 9, 12\}$
Clearly, every element of A is related to itself.
Therefore, it is a reflexive.
Now, R is not symmetry because 3 is related to 5 but 5 is not related to 3.
Also R is transitive relation because it satisfies the property that if aRb and bRc then aRc.

Question169

Let A = {1, 2, 3, 4} and R : A → A be the relation defined by R = {(1, 1), (2, 3), (3, 4), (4, 2)}. The correct statement is: [Online April 9, 2013]

Options:

A. R does not have an inverse.

B. R is not a one to one function.

C. R is an onto function.

D. R is not a function.

Answer: C

Solution:

Solution:
Domain = {1, 2, 3, 4}
Range = {1, 2, 3, 4}
∴ Domain = Range
Hence the relation R is onto function.

Question170

If P(S) denotes the set of all subsets of a given set S, then the number of one-to-one functions from the set S = {1, 2, 3} to the set P(S) is [Online May 19, 2012]

Options:

A. 24

B. 8

C. 336

D. 320

Answer: C

Solution:

Solution:
Let $S = \{1, 2, 3\} \rightarrow n(S) = 3$
Now, $P(S) =$ set of all subsets of S
total no. of subsets $= 2^3 = 8$
$\therefore n[P(S)] = 8$
Now, number of one-to-one functions from $S \rightarrow P(S)$ is ${}^8P_3 = \frac{8!}{5!} = 8 \times 7 \times 6 = 336$

Question171

If $A = \{x \in z^+ : x < 10$ and x is a multiple of 3 or 4$\}$, where z^+ is the set of positive integers, then the total number of symmetric relations on A is.
[Online May 12, 2012]

Options:

A. 2^5

B. 2^{15}

C. 2^{10}

D. 2^{20}

Answer: B

Solution:

Solution:
A relation on a set A is said to be symmetric iff (a, b) in $A \Rightarrow (b, a) \in A, \forall a, b \in A$
Here $A = \{3, 4, 6, 8, 9\}$
Number of order pairs of $A \times A = 5 \times 5 = 25$
Divide 25 order pairs of A times A in 3 parts as follows:
Part – A : (3, 3), (4, 4), (6, 6), (8, 8), (9, 9)
Part – B : (3, 4), (3, 6), (3, 8), (3, 9), (4, 6), (4, 8),(4, 9), (6, 8), (6, 9), (8, 9)
Part – C : (4, 3), (6, 3), (8, 3),(9, 3), (6, 4), (8, 4), (9, 4), (8, 6), (9, 6), (9, 8)
In part – A, both components of each order pair are same.
In part – B, both components are different but not two such order pairs are present in which first component of one order pair is the second component of another order pair and vice-versa.
In part–C, only reverse of the order pairs of part –B are present i.e., if (a, b) is present in part – B,

then (b, a) will be present in part –C
For example (3, 4) is present in part – B and (4, 3) present in part –C.
Number of order pair in A, B and C are 5, 10 and 10 respectively.
In any symmetric relation on set A, if any order pair of part –B is present then its reverse order pair of part –C will must be also present.
Hence number of symmetric relation on set A is equal to the number of all relations on a set D, which contains all the order pairs of part –A and part– B.
Now $n(D) = n(A) + n(B) = 5 + 10 = 15$
Hence number of all relations on set $D = (2)^{15}$
⇒ Number of symmetric relations on set $D = (2)^{15}$

Question172

The range of the function $f(x) = \frac{x}{1+|x|}$, $x \in R$, is [Online May 7, 2012]

Options:

A. R

B. (-1,1)

C. $R - \{0\}$

D. [-1,1]

Answer: B

Solution:

Solution:
$f(x) = \frac{x}{1+|x|}, x \in R$
If $x > 0, |x| = x \Rightarrow f(x) = \frac{x}{1+x}$
which is not defined for $x = -1$
If $x < 0, |x| = -x \Rightarrow f(x) = \frac{x}{1-x}$ which is not defined for $x = 1$
Thus $f(x)$ defined for all values of R except 1 and -1
Hence, range $= (-1, 1)$

Question173

Let A and B be non empty sets in R and $f : A \rightarrow B$ is a bijective function.
Statement 1: f is an onto function.
Statement 2: There exists a function $g : B \rightarrow A$ such that $fog = I_B$
[Online May 26, 2012]

Options:

A. Statement 1 is true, Statement 2 is false.

B. Statement 1 is true, Statement 2 is true; Statement 2 is a correct explanation for Statement 1.

C. Statement 1 is false, Statement 2 is true.

D. Statement 1 is true, Statement 2 is true, Statement 2 is not the correct explanation for Statement 1.

Answer: D

Solution:

Solution:
Let A and B be non-empty sets in R.
Let $f : A \rightarrow B$ is bijective function.
Clearly statement - 1 is true which says that f is an onto function.
Statement -2 is also true statement but it is not the correct explanation for statement-1

Question174

Let R be the set of real numbers.
Statement-1: $A = \{(x, y) \in R \times R : y - x$ is an integer $\}$ is an equivalence relation on R.
Statement- 2: $B = \{(x, y) \in R \times R : x = \alpha y$ for some rational number $\alpha\}$ is an equivalence relation on R.
[2011]

Options:

A. Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1.

B. Statement-1 is true, Statement-2 is false.

C. Statement-1 is false, Statement-2 is true.

D. Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.

Answer: A

Solution:

Solution:
$\because x - x = 0 \in I$ ($\therefore R$ is reflexive)
Let $(x, y) \in R$ as $x - y$ and $y - x \in I$ ($\because R$ is symmetric)
Now $x - y \in I$ and $y - z \in I \Rightarrow x - z \in I$
So, R is transative.
Hence R is equivalence.
Similarly as $x = \alpha y$ for $\alpha = 1$. B is reflexive symmetric and transative. Hence B is equivalence.
Both relations are equivalence but not the correct explanation.

Question175

The domain of the function $f(x) = \frac{1}{\sqrt{|x| - x}}$ is

[2011]

Options:

A. $(0, \infty)$

B. $(-\infty, 0)$

C. $(-\infty, \infty) - \{0\}$

D. $(-\infty, \infty)$

Answer: B

Solution:

Solution:
$f(x) = \frac{1}{\sqrt{|x| - x}}$, $f(x)$ is define if $|x| - x > 0$
$\Rightarrow |x| > x, \Rightarrow x < 0$
Hence domain of $f(x)$ is $(-\infty, 0)$

Question176

Let f be a function defined by
$f(x) = (x-1)^2 + 1, (x \geq 1)$
Statement -1: The set $\{x : f(x) = f^{-1}(x)\} = \{1, 2\}$
Statement -2: f is a bijection and $f^{-1}(x) = 1 + \sqrt{x-1}, x \geq 1$
[2011 RS]

Options:

A. Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.

B. Statement-1 is true, Statement-2 is true; Statement-2 is NOT a correct explanation for Statement-1.

C. Statement-1 is true, Statement-2 is false.

D. Statement-1 is false, Statement-2 is true.

Answer: A

Solution:

Solution:
Given f is a bijective function
$\therefore f : [1, \infty) \to [1, \infty)$
$f(x) = (x-1)^2 + 1, x \geq 1$
Let $y = (x-1)^2 + 1 \Rightarrow (x-1)^2 = y - 1$
$\Rightarrow x = 1 \pm \sqrt{y-1} \Rightarrow f^{-1}(y) = 1 \pm \sqrt{y-1}$
$\Rightarrow f^{-1}(x) = 1 + \sqrt{x-1} \{\therefore x \geq 1\}$
Hence statement- 2 is correct
Now $f(x) = f^{-1}(x)$
$\Rightarrow f(x) = x \Rightarrow (x-1)^2 + 1 = x$
$\Rightarrow x^2 - 3x + 2 = 0 \Rightarrow x = 1, 2$
Hence statement- 1 is correct

Question177

Consider the following relations:
R = { (x, y) | x, y are real numbers and x = wy for some rational

number w}; S = $\left\{ \left(\frac{m}{n}, \frac{p}{q} \right) \middle| \right.$ m, n, p. and q areintegers such that n, q ≠ 0 and qm = pn}
. Then
[2010]

Options:

A. Neither R nor S is an equivalence relation

B. S is an equivalence relation but R is not an equivalence relation

C. R and S both are equivalence relations

D. R is an equivalence relation but S is not an equivalence relation

Answer: B

Solution:

Solution:
Let xRy.
$\Rightarrow x = wy \Rightarrow y = \frac{x}{w}$
$\Rightarrow (y, x) \notin R$
R is not symmetric
Let S : $\frac{m}{n} S \frac{p}{q}$
$\Rightarrow qm = pn \Rightarrow \frac{p}{q} = \frac{m}{n}$
$\because \frac{m}{n} = \frac{m}{n} \therefore$ reflexive
$\frac{m}{n} = \frac{p}{q} \Rightarrow \frac{p}{q} = \frac{m}{n} \therefore$ symmetric
Let $\frac{m}{n} S \frac{p}{q}, \frac{p}{q} S \frac{r}{s}$
$\Rightarrow qm = pn, ps = rq$
$\Rightarrow \frac{p}{q} = \frac{m}{n} = \frac{r}{s}$
$\Rightarrow ms = rn$ transitive
.S is an equivalence relation.

Question178

Let $f(x) = (x+1)^2 - 1, x \geq -1$
Statement -1: The set $\{x : f(x) = f^{-1}(x) = \{0, -1\}$.

Statement- 2: f is a bijection.
[2009]

Options:

A. Statement-1 is true, Statement-2 is true. Statement-2 is not a correct explanation for Statement-1.

B. Statement-1 is true, Statement-2 is false.

C. Statement-1 is false, Statement-2 is true.

D. Statement-1 is true, Statement-2 is true. Statement-2 is a correct explanation for Statement-1.

Answer: D

Solution:

Solution:
Given that $f(x) = (x+1)^2 - 1, x \geq -1$
Clearly $D_f = [-1, \infty)$ but co-domain is not given. Therefore $f(x)$ is onto.
Let $f(x_1) = f(x_2)$
$\Rightarrow (x_1+1)^2 - 1 = (x_2+1)^2 - 1$
$\Rightarrow x_1 = x_2$
$\therefore f(x)$ is one-one, hence $f(x)$ is bijection
$\because (x+1)$ being something $+ve$, $\forall x > -1$
$\therefore f^{-1}(x)$ will exist. Let $(x+1)^2 - 1 = y$
$\Rightarrow x+1 = \sqrt{y+1}$ (+ve square root as $x+1 \geq 0$)
$\Rightarrow x = -1 + \sqrt{y+1}$
$\Rightarrow f^{-1}(x) = \sqrt{x+1} - 1$
Then $f(x) = f^{-1}(x)$
$\Rightarrow (x+1)^2 - 1 = \sqrt{x+1} - 1$
$\Rightarrow (x+1)^2 = \sqrt{x+1} \Rightarrow (x+1)^4 = (x+1)$
$\Rightarrow (x+1)[(x+1)^3 - 1] = 0 \Rightarrow x = -1, 0$
$\therefore$ The statement- 1 and statement- 2 both are true.

Question179

Let R be the real line. Consider the following subsets of the plane R × R :
S = {(x, y) : y = x + 1 and 0 < x < 2}
T = {(x, y) : x − y is an integer }

Which one of the following is true?
[2008]

Options:

A. Neither S nor T is an equivalence relation on R

B. Both S and T are equivalence relation on R

C. S is an equivalence relation on R but T is not

D. T is an equivalence relation on R but S is not

Answer: D

Solution:

Solution:
Given that
$S = \{(x, y) : y = x + 1$ and $0 < x < 2\}$
$\because x \neq x + 1$ for any $x \in (0, 2)$
$\Rightarrow (x, x) \notin S$
So, S is not reflexive.
Hence, S in not an equivalence relation.
Given $T = \{x, y) : x - y$ is an integer $\}$
$\because x - x = 0$ is an integer, $\forall x \in R$
So, T is reflexive.
Let $(x, y) \in T \Rightarrow x - y$ is an integer then $y - x$ is also an integer $\Rightarrow (y, x) \in R$
$\therefore T$ is symmetric
If $x - y$ is an integer and $y - z$ is an integer then $(x - y) + (y - z) = x - z$ is also an integer.
$\therefore T$ is transitive
Hence T is an equivalence relation.

Question180

Let $f : N \rightarrow Y$ be a function defined as $f(x) = 4x + 3$ where $Y = \{y \in N : y = 4x + 3$ for some $x \in N\}$.
Show that f is invertible and its inverse is
[2008]

Options:

A. $g(y) = \frac{3y + 4}{3}$

B. $g(y) = 4 + \frac{y+3}{4}$

C. $g(y) = \frac{y+3}{4}$

D. $g(y) = \frac{y-3}{4}$

Answer: D

Solution:

Solution:
Clearly $f(x) = 4x + 3$ is one one and onto, so it is invertible.
Let $f(x) = 4x + 3 = y$
$\Rightarrow x = \frac{y-3}{4}$
$\therefore g(y) = \frac{y-3}{4}$

Question181

Let W denote the words in the English dictionary. Definethe relation R by R = (x, y) ∈ W × W | the words x and yhave at least one letter in common.} Then R is
[2006]

Options:

A. not reflexive, symmetric and transitive

B. relexive, symmetric and not transitive

C. reflexive, symmetric and transitive

D. reflexive, not symmetric and transitive

Answer: B

Solution:

Solution:
Clearly $(x, x) \in R$, $\forall x \in W$
$\because$ All letter are common in some word. So R is reflexive.
Let $(x, y) \in R$, then $(y, x) \in R$ as x and y have at least one letter in common. So, R is symmetric.
But R is not transitive for example

Let x = BOY , y = TOY and z = THREE
then $(x, y) \in R$(O, Y are common) and $(y, z) \in R$(T is common) but $(x, z) \notin R$. (as no letter is common)

Question182

A real valued function f (x) satisfies the functional equation f (x − y) = f (x)f (y) − f (a − x)f (a + y) where a is a given constant and f (0) = 0, f (2a − x) is equal to
[2005]

Options:

A. $-f(x)$

B. $f(x)$

C. $f(a) + f(a - x)$

D. $f(-x)$

Answer: A

Solution:

Solution:
Given that $f(0) = 0$ and put
$x = 0, y = 0$
$f(0) = f^2(0) - f^2(a)$
$\Rightarrow f^2(a) = 0 \Rightarrow f(a) = 0$
$f(2a - x) = f(a - (x - a))$
$= f(a)f(x - a) - f(0)f(x)$
$= f(a)f(x - a) - f(x) = -f(x)$
$\Rightarrow f(2a - x) = -f(x)$

Question183

Let R = {(3, 3), (6, 6), (9, 9), (12, 12), (6, 12), (3, 9),(3, 12), (3, 6)} be a relation on the set A = {3, 6, 9, 12}. The relation is
[2005]

Options:

A. reflexive and transitive only

B. reflexive only

C. an equivalence relation

D. reflexive and symmetric only

Answer: A

Solution:

Solution:
R is reflexive and transitive only.
Here(3, 3), (6, 6), (9, 9), (12, 12) ∈ R [So, reflexive]
(3, 6), (6, 12), (3, 12) ∈ R[So, transitive]
∵(3, 6) ∈ R but (6,3)∉R[So, non-symmetric]

Question184

Let f : (−1, 1) → B, be a function defined byf (x) = $\tan^{-1}\frac{2x}{1-x^2}$, then f is both one − one and onto when B is the interval [2005]

Options:

A. $\left(0, \frac{\pi}{2}\right)$

B. $\left[0, \frac{\pi}{2}\right)$

C. $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

D. $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

Answer: D

Solution:

Solution:

Given $f(x) = \tan^{-1}\left(\frac{2x}{1-x^2}\right) = 2\tan^{-1}x$

for $x \in (-1, 1)$

If $x \in (-1, 1) \Rightarrow \tan^{-1}x \in \left(\frac{-\pi}{4}, \frac{\pi}{4}\right)$

$\Rightarrow 2\tan^{-1}x \in \left(\frac{-\pi}{2}, \frac{\pi}{2}\right)$

Clearly, range of $f(x) = \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

For f to be onto, codomain = range

$\therefore$ Co-domain of function $= B = \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

Question185

The graph of the function y = f (x) is symmetrical about the line x = 2, then
[2004]

Options:

A. $f(x) = -f(-x)$

B. $f(2 + x) = f(2 - x)$

C. $f(x) = f(-x)$

D. $f(x + 2) = f(x - 2)$

Answer: B

Solution:

Solution:
(b) Given that a graph symmetrical. with respect to line $x = 2$ as shown in the figure.



From the figure

$f(x_1) = f(x_2)$, where $x_1 = 2 - x$ and $x_2 = 2 + x$

$\therefore f(2 - x) = f(2 + x)$

Question186

Let R = {(1,3),(4, 2), (2, 4),(2,3),(3,1)} be a relation on the set A = {1, 2,3, 4}.. The relation R is
[2004]

Options:

A. reflexive

B. transitive

C. not symmetric

D. a function

Answer: C

Solution:

Solution:

$\because (1, 1) \notin R \Rightarrow R$ is not reflexive

$\because (2, 3) \in R$ but $(3,2) \notin R$

$\therefore R$ is not symmetric

$\because (4, 2)$ and $(2,4) \in R$ but $(4,4) \notin R$

$\Rightarrow R$ is not transitive

Question187

If f : R → S, defined by f (x) = sin x − $\sqrt{3}$ cos x + 1, is onto, then the interval of S is
[2004]

Options:

A. [-1,3]

B. [-1,1]

C. [0,1]

D. [0,3]

Answer: A

Solution:

Solution:
Given that $f(x)$ is onto
∴ range of $f(x)$ = codomain = S
Now, $f(x) = \sin x - \sqrt{3}\cos x + 1$

$= 2\sin\left(x - \frac{\pi}{3}\right) + 1$

we know that $-1 \le \sin\left(x - \frac{\pi}{3}\right) \le 1$

$-1 \le 2\sin\left(x - \frac{\pi}{3}\right) + 1 \le 3 \therefore f(x) \in [-1, 3] = S$

Question188

Domain of definition of the function f (x) = $\frac{3}{4 - x^2}$ + $\log_{10}(x^3 - x)$, is

[2003]

Options:

A. (-1,0)∪(1, 2) ∪ (2, ∞)

B. (a, 2)

C. (-1,0)∪(a, 2)

D. (1,2)∪(2, ∞)

Answer: A

Solution:

Solution:

$f(x) = \frac{3}{4-x^2} + \log_{10}(x^3 - x)$

$4 - x^2 \neq 0;\ x^3 - x > 0$

$x \neq \pm\sqrt{4}$ and $-1 < x < 0$ or $1 < x < \infty$



$\therefore D = (-1, 0) \cup (1, \infty) - \{\sqrt{4}\}$

$D = (-1, 0) \cup (1, 2) \cup (2, \infty)$

Question189

If $f : R \to R$ satisfies $f(x + y) = f(x) + f(y)$, for all $x, y \in R$ and $f(1) = 7$, then $\sum_{r=1}^{n} f(r)$ is

[2003]

Options:

A. $\frac{7n(n+1)}{2}$

B. $\frac{7n}{2}$

C. $\frac{7(n+1)}{2}$

D. $7n + (n + 1)$

Answer: A

Solution:

Solution:

$f(x + y) = f(x) + f(y)$

$\because f(1) = 7$

$f(2) = f(1 + 1) = f(1) + f(1) = 14$

$f(3) = f(1 + 2) = f(1) + f(2) = 21$

– – – – – – – – – – –

$$\therefore \sum_{r=1}^{n} f(r) = 7(1+2+3.....+n)$$
$$= \frac{7n(n+1)}{2}$$

Question190

A function f from the set of natural numbers to integers defined by

$$\mathbf{f(n) = } \begin{cases} \frac{n-1}{2}, \text{when n is odd} \\ -\frac{n}{2}, \text{when n is even} \end{cases} \mathbf{is}$$

[2003]

Options:

A. neither one -one nor onto

B. one-one but not onto

C. onto but not one-one

D. one-one and onto both.

Answer: D

Solution:

Solution:
We have $f : N \to I$
Let x and y are two even natural numbers, and $f(x) = f(y) \Rightarrow \frac{-x}{2} = \frac{-y}{2} \Rightarrow x = y$
$\therefore f(n)$ is one-one for even natural number.
Let x and y are two odd natural numbers and $f(x) = f(y) \Rightarrow \frac{x-1}{2} = \frac{y-1}{2} \Rightarrow x = y$
$\therefore f(n)$ is one-one for odd natural number.
Hence f is one-one.
Let $y = \frac{n-1}{2} \Rightarrow 2y + 1 = n$
This shows that n is always odd number for $y \in I$ .....(i)
and $y = \frac{-n}{2} \Rightarrow -2y = n$
This shows that n is always even number for $y \in I$ .....(ii)
From (i) and (ii)

Range of $f = I =$ codomain
$\therefore f$ is onto.
Hence f is one one and onto both.